

Cholecystokinin Antagonists: Pharmacological and Therapeutic Potential

Rosario Herranz

Instituto de Química Medica (CSIC), Juan de la Cierva 3, E-28006 Madrid, Spain

DOI 10.1002/med.10042



Abstract: Cholecystokinin (CCK) is a regulatory peptide hormone, predominantly found in the gastrointestinal tract, and a neurotransmitter present throughout the nervous system. In the gastrointestinal system CCK regulates motility, pancreatic enzyme secretion, gastric emptying, and gastric acid secretion. In the nervous system CCK is involved in anxiogenesis, satiety, nociception, and memory and learning processes. Moreover, CCK interacts with other neurotransmitters in some areas of the CNS. The biological effects of CCK are mediated by two specific G protein coupled receptor subtypes, termed CCK₁ and CCK₂. Over the past fifteen years the search of CCK receptor ligands has evolved from the initial CCK structure derived peptides towards peptidomimetic or non-peptide agonists and antagonists with improved pharmacokinetic profile. This research has provided a broad assortment of potent and selective CCK₁ and CCK₂ antagonists of diverse chemical structure. These antagonists have been discovered through optimization programs of lead compounds which were designed based on the structures of the C-terminal tetrapeptide, CCK-4, or the non-peptide natural compound, asperlicin, or derived from random screening programs. This review covers the main pharmacological and therapeutic aspects of these CCK₁ and CCK₂ antagonist. CCK₁ antagonists might have therapeutic potential for the treatment of pancreatic disorders and as prokinetics for the treatment of gastroesophageal reflux disease, bowel disorders, and gastroparesis. On the other hand, CCK₂ antagonists might have application for the treatment of gastric acid secretion and anxiety disorders. © 2003 Wiley Periodicals, Inc. *Med Res Rev*, 23, No. 5, 559–605, 2003

Key words: cholecystokinin; CCK₁ receptors; CCK₂ receptors; antagonists

1. INTRODUCTION

Cholecystokinin (CCK) was first discovered in 1928 as a hormone in the gastrointestinal tract,¹ which was identified in 1966 as a 33 amino acid peptide in porcine intestine extracts, and it was originally

Contract grant sponsor: CICYT; Contract grant number: SAF 2000-0147.

Correspondence to: Rosario Herranz, Instituto de Química Medica (CSIC), Juan de la Cierva 3, E-28006 Madrid, Spain.
E-mail: rosario@iqm.csic.es

named cholecystokinin-pancreozymin.^{2,3} Later, CCK was found in the central nervous system (CNS),⁴⁻⁷ and it is now generally believed to be the most widespread and abundant neuropeptide in the CNS. In the gastrointestinal tract CCK is released from endocrine cells in response to food intake, and regulates motility, contraction of gall bladder, pancreatic enzyme secretion, gastric emptying, and gastric acid secretion.⁸ In the nervous system CCK is involved in anxiogenesis,⁸⁻¹¹ satiety,^{8,12-14} nociception,^{8,15} thermoregulation,^{8,16} and memory and learning processes.^{9,17,18} Furthermore, the colocalization and interaction of CCK with other neurotransmitters in some CNS areas,^{8,19} mainly with dopamine (DA),^{20,21} suggests its implication in several neuropsychiatric disorders, such as schizophrenia, depression, and drug addiction.²⁰⁻²⁴

Besides the 33-amino acid sequence, CCK-33, formerly isolated and identified in porcine intestine, other species-specific molecular forms of CCK, derived from a 115-amino acid precursor protein (prepro-CCK²⁵), have been characterized later, such as CCK-58, CCK-39, CCK-22, CCK-8 [Asp-Tyr(SO₃H)-Met-Gly-Trp-Met-Asp-Phe-NH₂], unsulfated CCK-8, CCK-7, unsulfated CCK-7, CCK-5, and CCK-4 (Trp-Met-Asp-Phe-NH₂), which have in common the C-terminal sequence^{8,26} (Fig. 1). In humans CCK-58 and CCK-8 are the major forms.²⁷ This last octapeptide is relatively conserved across species, and appears to be the minimum sequence for full biological activity.²⁷

2. CCK RECEPTORS

The biological actions of CCK are mediated by two specific G protein coupled receptor (GPCR) subtypes, initially named CCK-A (for alimentary) and CCK-B (for brain).^{23,27} This nomenclature has been recently changed to CCK₁ and CCK₂, respectively, according to the guidelines of the International Union of Pharmacology (IUPHAR) Committee on Receptor Nomenclature and Drug Classification.²⁸ These receptors were pharmacologically classified on the basis of their affinity for the endogenous peptide CCK agonists and gastrin (Fig. 1), which share the same C-terminal pentapeptide amide sequence.

A. CCK₁ Receptors

As shown in Table I, these subtype receptors bind CCK-8 with a 500- to 1,000-fold higher affinity than gastrin or nonsulfated CCK-8, and a 10,000-fold higher affinity than CCK-4, and this binding is selectively inhibited by the antagonist devazepide (L-364,718).²⁷ These receptors are mainly localized in the gastrointestinal tract (pancreas, gall bladder, gastric mucosa, pyloric sphincter, sphincter of Oddi, and lower esophageal sphincter), where they are responsible of the regulation of




Figure 1. Amino acid sequence of the most predominant mammalian forms of CCK (CCK-58, CCK-33, CCK-8, and CCK-4, using single-letter amino acid symbols), and the major form of gastrin (gastrin-17). Filled circles show identical C-terminal pentapeptide sequence shared between CCK and gastrin. pQ, pyroglutamic acid.

Table 1. Summary of Characteristics, Localization, and Functions of CCK Receptor Subtypes

IUPHAR nomenclature ^a	CCK ₁	CCK ₂
Initial nomenclature ^a	CCK _A	CCK _B , CCK _B /gastrin
Endogenous agonists affinity rank order ^a	CCK-8 >> gastrin, unsulfated CCK-8 (500-1000-fold) > CCK-4 (10-fold)	CCK-8 > gastrin, unsulfated CCK-8 (0-10) > CCK-4 (60-fold)
Main selective antagonists in rank order of affinity (pKi) ^a	Devazepide (9.8), T-0632 (9.6), SR-27897 (9.2), IQM-95,333 (9.2), PD-140,548 (7.9-8.6), lorglumide (7.2)	YM-022 (10.2), L-740,093 (10.0), GV-150,013 (9.3), RP-73,870 (9.3), L-365,260 (7.5-8.7), LY-26,2691 (7.5)
G Protein coupling ^a	G _{q/11} , G _i ¹	G _{q/11}
Signal transduction ^b	PLC, IP ₃ , DAG, Ca ²⁺ PLA ₂ /arachidonic acid Camp	PLC, IP ₃ , DAG, Ca ²⁺
Gastrointestinal (GI) tract localization and functions ^b	<u>Pancreatic acini</u> : enzyme secretion, growth <u>Pancreatic islets</u> : insulin secretion <u>Gastric mucosa</u> : Pepsinogen release from chief cell Somatostatin release from D cells <u>Gallbladder and GI smooth muscle</u> : Contraction and motility	<u>Gastric mucosa</u> : Growth Gastric acid secretion from parietal cells Histamine release from ECL cells Pepsinogen release from chief cell Inhibition of somatostatin release in D cells
CNS and PNS localization and functions ^b	<u>Selected areas of the CNS and vagus nerve</u> : satiety <u>Posterior nucleus accumbens</u> : DA release <u>Dorsal horn of the spinal cord</u> : Antagonism of opiod analgesia	<u>Throughout CNS</u> : Anxiety, neuroprotection, memory and learning, drug addiction, Antagonism of analgesia <u>Dopaminergic pathways</u> : inhibition of DA release
Other localizations ^b	<u>Tumoral cell lines</u> : AR42J, CHP212	<u>Immune cells</u> : monocytes and T lymphocytes <u>Tumoral cells</u> : medullary thyroid, gastric and colonic carcinomas, AR42J, small cell lung carcinoma, astrocytomas, leiomyosarcoma

^aReference 55.

^bReferences 23 and 27.

diverse digestive processes.^{23,27,29} They are also present in select areas of the peripheral nervous system (vagus nerve), and the CNS [nucleus tractus solitarius (NTS), posterior nucleus accumbens, hypothalamic dorsomedial nucleus, dorsal horn of the spinal cord, and anterior pituitary corticotrophs],^{23,27} where they mediate the satiety effects of CCK,^{12,14} regulate an increase in dopamine release,^{20,27} and antagonize opiod analgesia.^{27,30,31} CCK₁ receptors have also been characterized in diverse tumoral cell lines, where they may mediate growth.^{27,32,33}

The CCK₁ receptor cDNA was first cloned from rat pancreas, which showed a 429 amino acid sequence,³⁴ subsequently it was cloned from guinea pig gall bladder, pancreas, and gastric chief cells,³⁵ human gall bladder,^{36,37} and rabbit stomach.³⁸ CCK₁ receptors are highly conserved among these species with an overall amino acid homology of 80%.²³ As above mentioned, these receptors belong to the superfamily of GPCR, whose activation starts the signal transduction cascade of phospholipase C (PLC), with the formation of the second messengers, inositol 1,4,5-triphosphate (IP₃) and 1,2-diacylglycerol (DAG), leading to the release of intracellular Ca²⁺ and the activation of protein kinase C (PKC).^{23,27} Depending on the agonist used, CCK₁ receptor activation may result in

stimulation of phospholipase A₂ (PLA₂), which regulates Ca²⁺ release through the arachidonic acid cascade.²³ Moreover, it has been shown that high concentrations of CCK-8 can stimulate the adenylate cyclase signal transduction pathway with formation of 3'5'-cyclic monophosphate (cAMP)^{23,27} (Table I).

B. CCK₂ Receptors

As summarized in Table I, the order of affinities for this receptor subtype is CCK-8 > gastrin or nonsulfated CCK-8 (0- to 10-fold) > CCK-4 (60-fold),²⁷ and the binding is selectively inhibited by the antagonist L-365,260. These receptors are mainly found throughout the CNS, where they have been proposed to be responsible of the anxiogenic,⁸⁻¹¹ and neuroprotective^{8,39-41} effects of CCK, and of the involvement of CCK in nociception,^{8,15,18} drug addiction,^{18,20-24} and memory and learning processes.^{18,23,27} In the dopaminergic pathways CCK₂ receptor activation inhibits dopamine release.^{8,20,27} In the gastrointestinal tract, CCK₂ receptors inhibit the release of somatostatin from D cells, and mediate the contraction of smooth muscle cells, the acid secretion from parietal cells, and the release of histamine from enterochromaffin-like (ECL) cells and pepsinogen from chief cells.^{23,27,29} These receptors are also present on pancreatic acinar cells in dogs and guinea pig, where they mediate growth but not enzyme secretion,²⁷ and on immune cells such as monocytes⁴² and T lymphocytes,⁴³ where their function is unknown. Like CCK₁, CCK₂ receptors are expressed in diverse tumors and tumor-derived cell lines such as medullary thyroid, gastric, colon, ovarian, and small cell lung carcinomas, astrocytomas, and certain pancreatic cell lines, where they elicit cellular proliferation.^{23,27,44}

CCK₂ receptors have been cloned from various sources:²³ rat brain and stomach, the pancreatic tumoral cell line AR4-2J, human brain and stomach, guinea pig gall bladder and stomach, brain and gastric enterochromaffin-like and parietal cells of *Mastomys natalensis*, calf pancreas, and a rabbit genomic library. The sequencing of the cDNA of the rat CCK₂ receptor showed the presence of 452 amino acids,⁴⁵ and this receptor subtype is highly conserved in humans, canine, guinea pig, calf, rabbit, *M. Natalensis*, and rat, with an overall amino acid identity of 72%.²³ These receptors were historically viewed as distinct of gastrin receptors on the basis of their different relative affinities for CCK and gastrin like peptides.⁴⁶ However, the cloning of human, guinea pig and rat,^{45,47} and canine²⁷ brain CCK₂ receptors resulted in a single cDNA identical to that for the canine parietal cell gastrin receptor.⁴⁸ More recently, additional CCK receptor subtypes could not be identify in the cloning of CCK₁ and CCK₂ receptors of guinea pig stomach smooth cells, by hybridization screening of a guinea pig smooth muscle cDNA library, using ³²P random primed labeled cDNA probes from cloned rat CCK₁ and CCK₂ receptor coding regions.⁴⁹ The previously observed differences in receptor pharmacology between CCK₂ and gastrin receptors have been attributed to tissue-specific differences in receptor processing or membrane lipids, variation in receptor preparations between tissues, and to interlaboratory variation.²⁷ On the other hand, studies on several tumoral cell lines have shown the existence of new gastrin and glycine-extended gastrin receptors, other than the CCK₂, whose activation has a trophic effect, although they have not been completely characterized.^{27,50-54}

In contrast to CCK₁, the signal-transduction cascade for CCK₂ receptors has been rather poorly characterized, in large part because of the difficulty of working with isolated neurons or isolated gastric mucosal cells expressing these receptors. Central CCK₂ receptors have not been proved to be linked to a well characterized second-messenger system in the brain, including the phosphoinositide system. However, in transfected cells (Cos, Chinese hamster ovary), it has been shown that like the CCK₁, the CCK₂ receptors couples to a pertussis toxin-insensitive G protein, causing activation of PLC, and subsequent formation of IP₃ and DAG, leading to the release of intracellular Ca²⁺ and the activation of PKC.^{23,27} (Table I).

CCK₁ and CCK₂ receptors share 48% amino acid identity,^{27,34,45} having the main amino acid sequence differences in the intra- and extracellular loops and in the outer third of the transmembrane

domains adjacent to the extracellular space.^{29,56–58} In spite of the high level of homology in the amino acid sequence for each CCK receptor subtype among species (72–80%), minor species-specific differences in receptor structure and distribution do occur that can result in significant pharmacological and physiological differences. In fact, mutagenesis studies in CCK receptors have shown that single amino acid substitutions may change ligand affinity^{59,60} and, even, the agonist/antagonist functionality,⁶¹ which in turn explain species-related differences. Therefore, it is important to consider the appropriate species for the intended experimental goals, and to be careful before extrapolating data from one animal species to another. Recently, several single nucleotide polymorphisms have been found in the gene structure of human cholecystokinin receptors, mainly in the CCK₂ subtype, which alter drug affinity and/or efficacy,⁶² and it has been postulated that these polymorphisms may influence the susceptibility to CCK-related diseases, such as schizophrenia,⁶³ Parkinson's disease,⁶⁴ and obesity.⁶⁵ On the other hand, in both CCK₁ and CCK₂ receptors, some ligands have shown functional heterogeneity. Thus, CCK dose-response studies using pancreatic acini typically reveal a biphasic dose-response relationship: stimulation at low CCK concentrations and inhibition at supramaximal concentrations, as it is the case of the CCK₁ receptor mediated amylase release, protein synthesis, and adenosinetriphosphatase activity.²⁷ Initially, this biphasic activity was explained due to the existence of two different binding sites for CCK₁ receptors. Later, as the cloning studies on CCK₁ and CCK₂ receptors have failed to identify additional members of the CCK receptor family, their functional heterogeneity has been explained on the basis of the existence of different interconverting conformational states, with preferential or differential G protein coupling, and, therefore, distinct biological signal transduction and target cell function.^{18,27} Studies with radiolabeled ligands, mainly by comparing results obtained from binding of agonists and antagonist, have shown that CCK₁ receptors exist in three different affinity states: the agonist [¹²⁵I]CCK-8 identified the high- and the low-affinity states, while the antagonist devazepide bound to the low-affinity state and to the very-low-affinity state, which represent 80% of the receptors.^{18,66} Recently, it has been shown that, under physiological conditions, CCK acts on high- and low-affinity CCK₁ receptors present on distinct vagal afferent fibers.⁶⁷ CCK₂ receptors have also been shown to exist in three different affinity states.¹⁸ Competition radioligand binding studies with diverse CCK₂ receptor antagonists have provided evidence for the existence of two binding sites, or binding states, for this receptor subtype in gastric glands and cortex of some animal species.^{68–70} Furthermore, the distinct behavioral responses elicited by the CCK₂ selective agonists BC 197 (anxiogenic) and CB 264 (non-anxiogenic and increases memory and attention) have been also related with the existence of two different binding states of these receptors in the brain.^{71,72}

3. CCK RECEPTOR ANTAGONISTS

The variety of physiological effects of CCK and its possible role in some pathological disorders have stimulated research in this area and, over the past decade, a number of potent and selective CCK₁ and CCK₂ receptor agonists and antagonists have been reported, which have highly contributed, as useful pharmacological tools, to the characterization of CCK receptor subtypes and for gaining insight into the functional significance of CCK in the periphery and in the CNS. The high therapeutic potentiality of some of these CCK receptor ligands have led them to clinical studies, and a few of them are considered very promising drug candidates. Thus, CCK₁ agonists have been studied as satiety agents in the treatment of obesity, while CCK₁ antagonists have been studied for the treatment of pancreatic and functional bowel disorders, gastroesophageal reflux, and pancreatic cancer. On the other hand, CCK₂ antagonists have been studied for the control of anxiety and gastric acid secretion, and for the treatment of drug tolerance. However, the involvement of CCK₂ receptor activation in gastric acid secretion and in the anxiogenic effects of CCK has discouraged the clinical studies of agonists for these receptor subtype. Several reviews covering diverse aspects of CCK receptor agonists and

antagonists have been published over the last years.^{73–83} This article reviews the main advances in the development of CCK receptor antagonists, focusing mainly on those that have shown a greater pharmacological potential and those that could have a more promising therapeutic prospect.

Initially, most of the first CCK receptor antagonists were peptides or pseudopeptides derived from the structural modification of the amino acid sequence of CCK-7 and CCK-4. The *N*-Boc derivative of the C-terminal tripeptide, Boc-Met-Asp-Phe-NH₂, which inhibited the CCK-8-stimulated amylase release from guinea pig pancreatic acini with an IC₅₀ of 0.25 mM,⁸⁴ was the first of these peptidic antagonists. Deletion of the C-terminal Phe³³ residue from Boc-CCK-4⁸⁵ and Boc-CCK-7^{86,87} derivatives produced CCK₂ and CCK₁ antagonists with binding affinities in the 10^{–6} M range. Replacement of the Met³¹ residue in these tri- and hexapeptide derivatives by Orn(Z) led to the CCK₁ receptor antagonists Boc-Trp³⁰-Orn(Z)³¹-Asp³²-NH₂ and Boc-Tyr²⁷(SO₃H)-Nle²⁸-Gly²⁹-Trp³⁰-Orn(Z)³¹-Asp³²-NH₂, which showed binding potencies in the 10^{–7} M range.⁸⁸ It is interesting to note that, as the replacement of the Met residues of CCK-4 and CCK-8 derivatives by Nle or Leu has no influence on their biological activity. Therefore, most of the synthetic CCK derivatives include this replacement to avoid the Met instability. The combined replacement of Phe³³ by 2-phenylethyl ester⁸⁹ or amide,⁹⁰ and Trp³⁰ by D-Trp in Boc-CCK-7 produced the most potent peptidic CCK₁ receptor antagonists with binding affinities at these receptors in the 10^{–8} range. On the other hand, the replacement of the Met³¹ and Phe³³ residues of Boc-CCK-4 by the unnatural and hydrophobic amino acids phenylglycine (Phg) and dimethylamide-1-naphtylalanine [1-Nal-*N*(CH₃)₂], to increase metabolic stability and brain penetration, led to a CCK₂ antagonist with a binding affinity at CCK₂ receptors of 3.9 × 10^{–8} M.⁹¹ Finally, within the pseudopeptide derivatives, replacement of the peptide bonds Nle³¹-Asp³² in Boc-[Nle³¹]-CCK-4 and Trp³⁰-Nle³¹ in Z-[Nle³¹]-CCK-4 by the peptide bond surrogates ψ[CH₂NH]⁹² and Ψ[CH(CN)NH],⁹³ led to CCK₂ antagonists, which displayed binding affinities in the range of 10^{–7} and 10^{–8} M, respectively. Similarly, the replacement of the Nle²⁹-Gly²⁹ peptide bond in Boc-[Nle^{28,31}]-CCK-7 by the peptide bond surrogates Ψ[COCH₂] or Ψ[HNCO] gave the most potent peptidic CCK₂ antagonists, which displayed binding affinities in the subnanomolar range.⁹⁴ However, despite the very good potency and selectivity of some of these peptidic antagonists, their poor oral availability, low metabolic stability, difficulty to cross the blood-brain barrier, and, in some cases, their mixed antagonist/agonist character have hampered their therapeutic development. To avoid these problems, the search of CCK receptor ligands, both antagonists and agonists, has evolved towards the search of peptidomimetics or non-peptide ligands. The discovery in 1985 of the natural non-peptide compound asperlicin (**1**)⁹⁵ as a moderate, competitive, and selective antagonist of CCK₁ receptors (IC₅₀ = 10^{–6} M, pancreas binding)⁹⁶ represented a key breakthrough in this search. This finding opened the door to the search of peptidomimetics through random screening, not only in the CCK field, but in general in most peptide receptors. As it will be discussed later, the structure manipulation of asperlicin has led to the discovery of benzodiazepine and the quinazolidinone derivatives (Fig. 2) as two families of potent and selective CCK receptor antagonists. Over the past decade, antagonists with high structural diversity have emerged, which will be commented below grouped in structure families. In most of these groups, there are selective antagonists for CCK₁ and CCK₂ receptors, which will be discussed separately according to their selectivity.

A. Amino Acid Derivatives

1. CCK₁ Receptor Antagonists

Benzoyl-DL-glutamic acid dipropylamide, named proglumide (Table II, compound **3**), reported in 1967 by researchers at Rotta Laboratories as a weak gastrin antagonist,⁹⁷ could be considered as the first discovered CCK receptor antagonist. This compound, along with *p*-chlorobenzoyl-L-tryptophan (benzotript, **2**), was tested for its ability to antagonize the effects of CCK in 1981.^{98,99} Both





Figure 2. General structures of 1,4-benzodiazepine and quinazolidinone derivatives, potent and selective CCK receptor antagonist, derived from manipulation of the asperlicin structure.

compounds were found to be weak competitive non selective antagonists of comparable potency in the inhibition of the [¹²⁵I]CCK-8 binding, the gastrin-stimulated gastric acid secretion, and CCK-stimulated amylase release from pancreatic acini. Proglumide has been marketed by Rotta Laboratories in Europe for the treatment of ulcers. Additional derivatives of tryptophan,¹⁰⁰ as well as simple analogues of most of the other coded amino acids,¹⁰¹ have been investigated as potential CCK antagonists. However, it has been the group of the glutamic acid derivatives which, mainly by **manipulation of the aromatic *N*-acyl group and the *N*-alkyl-carboxamide at the α -carboxy group**, has provided the most potent and selective antagonists for both CCK₁ and CCK₂ receptors. To illustrate the rank order of potency, some of these compounds are summarized in Tables II and III with their binding affinity data. It must be noted that is not possible to do a precise quantitative comparison with data reported for different compounds, as they were issued from different research teams, sometimes using different animal species, tests, or experimental conditions. Therefore, these comparisons must be taken with precaution. On the benzoyl group of proglumide both steric and electronic effects, as well as regiochemistry, were explored with a variety of substituents.¹⁰² The 3,4-dichloro substitution was which gave the most potent compounds. The alkyl amide groups were also found to play an important role. Thus, enlargement of the propyl groups of proglumide to *n*-pentyl, along with the 3,4-dichloro substitution at the benzoyl group, led to a higher than 10⁴ increase in the CCK₁ binding affinity of lorglumide (4, CR 1409) a moderate and very selective CCK₁ antagonist, which displayed activity after oral administration and low toxicity.¹⁰³ As shown in Table II, the resolved *D*-enantiomer of this glutamic acid derivative is more potent at CCK₁ receptors than the corresponding *L*-enantiomer (compounds 5 and 6). Loxiglumide (7, CR 1505), analogue of lorglumide in which an oxygen replaces one methylene, has shown very similar antagonistic properties to those of lorglumide.^{104,105} Both compounds are moderately potent, competitive and selective CCK₁ receptor antagonists *in vitro* and *in vivo*, and are devoid of agonist activity.

In spite of the moderate potency of loxiglumide, its good pharmacokinetic properties and oral bioavailability have facilitated its use in pharmacological and clinical studies,¹⁰⁶ having been one of the CCK₁ antagonists more widely studied. Studies in animals have shown that this antagonist protects against experimental pancreatitis,¹⁰⁷ and these results were the basis for

Table II. Most Significant Selective CCK₁ Receptor Antagonists Derived From Amino Acids

Compound	Config. (*)	R ¹	R ²	R ³	IC ₅₀ (μM)		Selectivity CCK ₂ /CCK ₁
					CCK ₁	CCK ₂	
Benzotript ^a (2)					82	40	0.49
Proglumide ^a (3)	DL	<i>n</i> -Pr	<i>n</i> -Pr		6,100	11,000	1.8
Lorglumide ^a (4) (CR 1409)	DL	<i>n</i> -Pentyl	<i>n</i> -Pentyl		0.13	300	2,307
D-Lorglumide ^a (5)	D	<i>n</i> -Pentyl	<i>n</i> -Pentyl		0.07	NR ^b	
L-Lorglumide ^a (6)	L	<i>n</i> -Pentyl	<i>n</i> -Pentyl		5.1	NR ^b	
Loxiglumide ^c (7) (CR 1505)	DL	CH ₃ O(CH ₂) ₃	<i>n</i> -Pentyl		0.33	9.1	27
Dexloxiglumide ^c (8) (CR 2017)	D	CH ₃ O(CH ₂) ₃	<i>n</i> -Pentyl		0.12	22	170
A-64718 ^d (9)	D	<i>n</i> -Pentyl	<i>n</i> -Pentyl		0.02	1.30	68
A-65186 ^d (10)	D	<i>n</i> -Pentyl	<i>n</i> -Pentyl		0.005	3.50	686

^aReference 103. Binding affinities in rat pancreatic acini (CCK₁) and mouse brain cortex (CCK₂), using the radioligand [¹²⁵I]CCK-8.


^bNR = non reported.

^cReference 104.

^dReference 115. Binding affinities in guinea pig pancreas and brain, using the radioligand [¹²⁵I]CCK-8.

its clinical evaluation for the treatment of acute pancreatitis.¹⁰⁸ In healthy subjects loxiglumide inhibits the slow, but not the rapid phase of postprandial gall bladder contraction and accelerates gastric emptying.^{109,110} It does not affect gastric motility or sensitivity during duodenal saline infusion, but partially restores gastric tonic activity during lipid infusion, reduces the occurrence of meal-like fullness and nausea, and increases the pressure at which sensations are reported.¹¹¹ Loxiglumide also inhibits the transient lower oesophageal sphincter relaxations and attenuates the fall in lower oesophageal sphincter (LES) pressure following a meal.¹¹² The injectable formulation of loxiglumide, for the treatment of acute pancreatitis, has been submitted for regulatory approval in Japan, and its oral formulation is undergoing Phase III trials for chronic pancreatitis.

Dexloxiglumide (8, CR 2017), retains all pharmacological properties of the racemic compound loxiglumide, but is more potent than this or its L-enantiomer. As loxiglumide, dexloxiglumide

Table III. Most Significant Selective CCK₂ Receptor Antagonists Derived From Amino Acids

Compound	R	IC ₅₀ (nM)		Selectivity CCK ₁ /CCK ₂
		CCK ₁	CCK ₂	
Spiroglumide ^a (13) (CR 2194)	OH	13,500	1,400	9.6
CR 2345 ^b (14)		6,600	700	9.4
CR 2767 ^a (15)		3,400	57	60
CR 2622 ^a (16)		7,380	20	369
 17 Itriglumide ^c (CR 2945)		20,700	2.3	9,000

^aData reported in reference 125. Receptor binding affinities determined in rat pancreatic acini, using the radioligand [¹²⁵I]CCK-8 (CCK₁), and in guinea pig cortex, using the radioligand [³H]-N-Me-N-Leu-CCK-8 (CCK₂).

^bData reported in reference 78. Receptor binding affinities determined as for **13**.

^cData reported in reference 127. Receptor binding affinities determined in rat pancreatic acini, using the radioligand [¹²⁵I]CCK-8 (CCK₁), and in rat cortex, using the radioligand [³H]pBC264 (CCK₂).

has a good pharmacokinetic profile, and it has shown good safety and tolerability.¹¹³ Results from both preclinical and clinical studies with this CCK₁ antagonist indicate that it is an effective inhibitor of gall bladder contraction, improves lower oesophageal sphincter function, and accelerates gastric emptying and colonic transit. Dexloxiglumide also significantly decreases symptoms in irritable bowel syndrome (IBS) and functional dyspepsia patients. Therefore, has potential as an effective treatment for constipation-predominant IBS, functional dyspepsia, constipation, LES function, gastric emptying disorders and biliary colics. Phase III studies with dexloxiglumide are in progress in USA, and Forest Laboratories has entered into an agreement with Rotta for its development and marketing. In June 2001, Merrill Lynch predicted an USA filing date for this compound in 2003.¹¹⁴

More potent CCK₁ antagonists have been discovered combining features of lorglumide and benzodiazepine structures,¹¹⁵⁻¹¹⁷ such as A-64718 (**9**) and A-65186 (**10**), which showed nanomolar




Figure 3.

affinities at CCK₁ receptors, although data of pharmacological or clinical studies have not been reported. Additionally, the analogue of loxiglumide KSG-504¹¹⁸ (Fig. 3, **11**), and the aspartic acid derivative 2-NAP^{119,120} (**12**), both incorporating a 2-naphthalenesulphonyl group, have been described as competitive and selective CCK₁ antagonists, with binding affinities in the 10⁻⁷ M range, and they have been studied in Phase I clinical trials.

2. CCK₂ Receptor Antagonists

The structure modification of lorglumide also produced selective CCK₂ receptor antagonists.¹²¹ Thus, spiroglumide (Table III, compound **13**) showed competitive and specific antagonism of the pentagastrin-stimulated gastric acid secretion in several animal species and models.¹²² In humans, this CCK₂ antagonist dose-dependently antagonized gastrin-stimulated gastric acid and fluid responses with a competitive-like profile.¹²³ In a subsequent Phase I randomized, double-blind, placebo-controlled trial in healthy male volunteers, spiroglumide significantly decreased basal acid output in response to food ingestion as well as postprandial intragastric acidity.¹²⁴ However, despite its excellent oral bioavailability, the low affinity and selectivity at CCK₂ receptors (micromolar range) precluded further development of spiroglumide as a potential therapeutic tool for peptic ulcer diseases.⁷⁸

The chemical manipulation of the structure of spiroglumide has led to more potent and selective CCK₂ antagonists.¹²⁵ Among these, the *N*-methylpiperazinyl derivative CR 2345 (**14**) was synthesized to investigate whether the acidic character of glutamic acid derivatives was essential for CCK₂ antagonism. The basic substitution in the carboxylic acid with the methylpiperazinyl group fully retains antagonistic activity. Moreover, CR 2345 also inhibited histamine- and carbachol-induced acid secretion.¹²⁶ Interestingly, this compound, as well as its *N*-propyl analogue (CR 2456), was shown to inhibit the growth *in vitro* of several human strains of *Helicobacter pylori*, isolated from patients with histological lesions of gastritis, now recognized as the aetiological factor of peptic ulcers, although the MICs of these spiroglumide derivatives were high (1–16 µg/ml) in comparison to those of the most effective antibiotics (e.g., the MIC₅₀ range for clarithromycin is 0.03–0.3 µg/ml).⁷⁸ More potent and selective spiroglumide derivatives were obtained when the amino acids *N*-methyl-L-tryptophan (**15**, CR 2767) or L-glutamic acid *N*-(1-naphthyl)-amide (**16**, CR 2622) were coupled to the side chain carboxylic acid of spiroglumide.¹²⁵ Compound **16** given intravenously reduced dose-dependently the pentagastrin-stimulated gastric acid secretion in rat stomach. However, the relatively high molecular weight of this compound limits its oral absorption.⁷⁸ Further structure manipulation of CR 2622 has led to the anthranilic acid derivative itriglumide (**17**, CR 2945). This lower molecular weight analogue has improved *in vivo* oral bioavailability, and showed nanomolar affinity and excellent CCK₂ selectivity (CCK₁/CCK₂ = 9,000).¹²⁷ *In vivo*, itriglumide blocked pentagastrin-induced gastric acid secretion in anaesthetized rats with a 50% inhibitory dose of 0.3 mg/kg given intravenously, and 4.0 mg/kg given intraduodenum. In this type of administration, it was more potent than both ranitidine and omeprazole. The gastrin antagonism was reversible and competitive, with a pA₂ value of 7.33, and was gastrin-specific, as it was unable to antagonize the gastric acid secretion

stimulated by histamine or carbachol.¹²⁸ This compound also showed to be effective in prevention of gastric damage in several models.⁷⁸ Furthermore, itriglumide showed significant dose-dependent anxiolytic-like effects in four rodent tests of anxiety, comparable to those of the anxiolytic diazepam, but without signs of sedation and ataxia. Additionally, a 7-day repeated treatment with this compound at 10 mg/kg/day s.c. did not induced tolerance or withdrawal anxiety in rats.¹²⁸ Itriglumide also displayed antiproliferative effects in mice GI tumors.¹²⁹ In healthy volunteers, this CCK₂ antagonist was well tolerated, and now is in Phase I clinical studies as anxiolytic and antiulcer.

B. 1,4-Benzodiazepine Derivatives


1. CCK₁ Receptor Antagonists

As it has been mentioned above, the chemical manipulation of asperlicin (**1**) structure, retaining the 1,4-benzodiazepine skeleton, has given very potent and selective antagonists for both CCK receptor subtypes CCK₁ and CCK₂. The first discoveries in this family of antagonists were made by Merck chemists, reasoning that combining the elements of diazepam with D-tryptophan might mimic asperlicin in its CCK receptor affinity. This idea proved to be very successful, obtaining the first non-peptide selective CCK₁ antagonists, which displayed similar affinity to that of asperlicin and oral activity.^{130,131} Efforts to optimize the CCK₁ activity of these first benzodiazepine derivatives yielded the highly potent and selective antagonist devazepide (Table IV, compound **18**, also known as L-364,718 or MK-329).^{132,133} This antagonist displays subnanomolar affinity at CCK₁ receptors, comparable to that of the native ligand CCK-8, and high selectivity versus the CCK₂. Devazepide has been shown to be highly potent by several routes of administration, including oral, in a variety of functional assays such as gastric emptying and gall bladder contraction in a number of different species, and no agonistic activity has been observed.⁷³ It has also been demonstrated that devazepide crosses the blood-brain barrier efficiently.^{134,135} Due to its high potency, selectivity and bioavailability, devazepide has been the reference CCK₁ receptor antagonist more widely used as pharmacological tool for the study of the physiological effects of CCK and its receptors. In Phase I clinical trials, in healthy human volunteers, devazepide inhibited CCK-induced gall bladder contraction,^{136,137} and stimulated gastric motility and gastric emptying after ingestion of meals, although this stimulation depends on the meal lipid content.¹³⁸ Although no data have been reported on Phase II clinical trials, Merck discontinued the development of devazepide in this phase in 2001, due to gallstone toxicity,^{76,139} and it has been reported that this compound induces hyperplasia in the rat liver and bile ducts.¹⁴⁰ In spite of this, ML Laboratories and Panos Therapeutics have recently initiated a joint full Phase II clinical study of devazepide for the treatment of pain, in patients suffering from opioid-resistant neuropathic pain.

All the chemical modifications upon the devazepide structure have been detrimental both for the binding potency and for the selectivity at CCK₁ receptors. The inversion of the absolute configuration at the C-3 of the 1,4-benzodiazepine skeleton led to a two order of magnitude decrease in the CCK₁ potency for the enantiomer (*R*)-devazepide (Table IV, **19**). As it will be shown below, the decrease in CCK₁ selectivity with this inversion of configuration at C-3 is general in the 1,4-benzodiazepine-derived antagonists, and in most of them that configuration inversion produced a reversal of selectivity, leading to CCK₂ selective antagonists. Among the devazepide analogues developed as CCK₁ antagonists, pranazepide (FK-480, **20**)^{141,142} and tarazepide (**21**)^{143,144} have reached Phase II clinical trials, although no data are available in the peer-reviewed literature about the results of these compounds in human clinical studies.

2. CCK₂ Receptor Antagonists

Researchers at Merck also discovered the first highly potent and selective non-peptide CCK₂ receptor antagonist L-365,260 (Table V, compound **22**) in the process of asperlicin structure manipulation,

Table IV. Most Significant 1,4-Benzodiazepine-Derived Selective CCK₁ Receptor Antagonists


Compound	IC ₅₀ (nM)		Selectivity CCK ₂ /CCK ₁	Reference
	CCK ₁ ^a	CCK ₂ ^b		
Asperlicin (1)	1,400	>10 ⁵	>71	130
Devazepide (18) (L-364,718 or MK-329)	0.08	270	3,375	133
(<i>R</i>)-Devazepide (19)	8.3	3,700	446	133
Pranazepide (20) (FK-480)	0.67	310	463	141
Tarazepide (21)	NR ^c	NR ^c		143

^aBinding affinities in rat pancreas, using the radioligand [¹²⁵I]CCK-33 except for **20** that [¹²⁵I]CCK-8 was used.

^bBinding affinities in guinea pig brain, using the radioligand [¹²⁵I]CCK-33 except for **20** that [¹²⁵I]CCK-8 was used.

^cData not reported.

through the combination of replacing the indol-2-yl-amide of devazepide by an aryl urea moiety at the 1,4-benzodiazepine C-3 position and the (*R*) stereochemistry at that position.¹⁴⁵ That compound, which displayed nanomolar affinity at CCK₂ receptors and 140-fold selectivity versus the CCK₁, also represented a significant landmark in the pharmacological studies of CCK and its receptors. Upon oral administration, it potently antagonized gastrin-stimulated acid secretion in several animal species with good duration of action.¹⁴⁶ However, in Phase I clinical trials in healthy male volunteers, L-365,260 produced only a modest inhibition of gastrin-stimulated gastric acid secretion, and this effect was of short duration.¹⁴⁷ In spite of the predominant role of CCK₂ receptors in the anxiogenic effects of CCK, L-365,260 has not clearly showed anxiolytic activity in animal models.¹⁰ In Phase II clinical studies, this antagonist achieved a significant reduction in the frequency and intensity of CCK-4¹⁴⁸ and lactate-induced¹⁴⁹ panic attacks in patients with panic disorders, and has shown to reverse the autonomic and anxiogenic effects of pentagastrin in healthy volunteers.¹⁵⁰ However, in another study, it was ineffective in limiting panic attacks in patients.¹⁵¹ These discouraging clinical results were attributed to its limited oral bioavailability, which could be explained by its low aqueous solubility.^{152,153} Based on the modest results of L-365,260 in the inhibition of gastric acid secretion, Merck researchers envisaged only a marginal role in the antiulcer therapy for this CCK₂ antagonist, and searched for other analogues with greater CNS penetration and oral bioavailability with the aim of successfully controlling anxiety and panic disorders. This goal was pursued through the incorporation of bulkier substituents at the N-1 position and water-solubilizing groups into the structure of L-365,260. Thus,

Table V. Most Significant 1,4-Benzodiazepine-Derived Selective CCK₂ Receptor Antagonists

Compound	R ¹	R ²	R ³	IC ₅₀ (nM)		Selectivity CCK ₁ /CCK ₂	Reference
				CCK ₁ ^a	CCK ₂ ^b		
L-365,260 (22)	CH ₃	Phenyl	CH ₃	280	2.0	140	145
23 ^c	CH ₂ CON(Et) ₂	Phenyl	OCH ₃	120	0.22	545	154
L-708,474 (24)	CH ₃	Cyclohexyl	CH ₃	1,797	0.28	6,418	155
L-368,730 (25)	CH ₃	Phenyl		577	1.0	577	156
L-369,466 (26)	CH ₃	Phenyl		983	0.27	3,640	156
L-736,380 (27)	CH ₃	Cyclohexyl		400	0.05	8,000	157
28 ^c	CH ₃		CH ₃	65	66	1	154
L-740,093 (29)	CH ₃		CH ₃	1604	0.10	16,040	161
YM022 (30)		Phenyl	CH ₃	150 ^d	0.11 ^e	1,364	163
YF476 (31)	CH ₂ COC(CH ₃) ₃		NHCH ₃	502 ^d	0.11 ^e	5,020	168

^aInhibition of the binding of [¹²⁵I]CCK-8 to rat pancreas tissues.^bInhibition of the binding of [¹²⁵I]CCK-8 to guinea pig cerebral cortex.^c(3*RS*)-Stereochemistry.^dInhibition of the binding of [³H]L-364,718 to rat pancreas membranes.^eInhibition of the binding of [¹²⁵I]CCK-8 to rat cerebral cortex.

more potent analogues were obtained either by the introduction of a diethyl acetamide group at N-1 as in **23**¹⁵⁴ (Table V), or by saturation of the 5-phenyl substituent to the cyclohexyl in L-708,474¹⁵⁵ (**24**). However, the water solubility of these compounds was also very low. In the second-generation benzodiazepines this issue was resolved, whilst maintaining the increased receptor affinity, firstly by replacing the 3-methyl substituent on the aryl urea moiety by acidic solubilizing groups such as the tetrazol or 1,2,4-oxadiazol-2-one rings.¹⁵⁶ Prominent compounds emerging from this series include the tetrazole derivatives L-368,730 (**25**) and L-368,935, and the oxadiazolone analogue L-369,466 (**26**). However, despite the excellent potency and increased solubility, these compounds did not exhibit satisfactory bioavailability. Therefore, further modifications were performed on the tetrazole moiety of L-368,730 in order to influence bioavailability by modulating the p*K*_A of the acidic moiety by means of structural manipulations around the tetrazole group.¹⁵⁷ From this study, the amino-tetrazole group was chosen as acidic moiety, because its p*K*_A (6.0) is substantially higher than that of the tetrazole itself. Compounds resulting from this strategy, such as L-736,380 (**27**) and its analogue L-738,425, in which a 2-tetrazolyl-isoindolyl moiety replaces the 3-(*N*-methyl-*N*-tetrazolyl)amino-phenyl of **27**, are among the most potent and, in the case of L-738,425 [CCK₂ IC₅₀ = 0.11 nM, CCK₁/CCK₂ = 37,000], most selective CCK₂ antagonists so far reported. L-736,380, given by the

intraperitoneal route, dose-dependently inhibited the gastric acid secretion in anaesthetized rats, with an ID_{50} of 0.064 mg/kg. However, these acidic compounds are significantly less brain penetrant than the prototype L-365,260 (22).¹⁵⁷

Only with the introduction of cationic water solubilizing groups, particularly within the C-5 substituent, the oral bioavailability was improved. Such strategy was initially prompted by the 4-fold increase in plasma concentration after oral dosing in rats, with respect to L-365,260, achieved in the case of the 2-pyridyl substituted derivative 28.¹⁵⁴ This change was accompanied by a reduction in receptor affinity and selectivity, and although this was restored by the introduction of bulkier substituents into the N-1 position, there was a concomitant reduction in aqueous solubility from this change, which was reflected in decreased plasma concentration with respect to L-365,260 upon oral administration. The optimization of this strategy has provided several improved compounds,^{158–160} standing out among them L-740,093 (29), where the introduction of the bulky basic 2-azabicyclo[3.2.2]non-2-yl moiety into C-5 increased both receptor affinity and aqueous solubility.¹⁶¹ According to its *in vitro* affinity, log D (4.7), aqueous solubility (0.15 mg/ml at pH = 5.0), and pK_A (7.1), L-740,093 was 100-fold more potent than L-365,260 in inhibition of pentagastrin-stimulated acid secretion upon i.p. administration.¹⁶¹

Although the second generation of 1,4-benzodiazepine-based CCK_2 receptor antagonists have superseded L-365,260 in terms of their *in vivo* properties in animal models, corresponding data in humans have so far not been reported. The recognition by Merck that L-365,260 and related 1,4-benzodiazepine-based CCK_2 receptor antagonists blocked the slowly activating component (I_{Ks}) of delayed rectifier potassium currents in guinea pig myocytes,¹⁶² and the association of this effect with cardiac arrhythmia, may have restricted subsequent clinical studies.

Yamanouchi in collaboration with Ferring developed the finding that CCK_2 receptor affinity of 1,4-benzodiazepine-based antagonists increased on introducing bulky substituents at the N-1 position. The optimal compound of this series, YM022 (30), which contains a 2-methyl-benzoyl-methyl group at this position, showed subnanomolar affinity at rat brain CCK_2 receptors, more than 2 orders of magnitude higher than that for rat pancreatic $CCK-A$ receptor.¹⁶³ *In vivo*, this compound inhibits gastrin-stimulated acid secretion in anaesthetized rats by i.v. administration (ED_{50} = 7.8 nM/kg).¹⁶⁴ Although YM022, as the other 1,4-benzodiazepines which contain bulky substituents at N-1, suffers from low aqueous solubility and requires formulation as a solid dispersion to achieve adequate oral bioavailability,¹⁶⁵ by this route was as effective as the histamine H_2 -antagonist famotidine and 8-fold more potent than the prototypical proton pump inhibitor omeprazole in models of gastric and duodenal ulcer in rats. YM022 had no effect on histamine or bethanechol-stimulated gastric acid secretion.¹⁶⁶ Moreover, YM022, given orally, was able to practically inhibit gastric damage induced by restraint stress, and inhibited the hypersecretion observed following cessation of omeprazole treatment.¹⁶⁷ These results supported the assertion that this compound may have a role as an alternative anti-ulcer therapy devoid of the risk in relapse, and promoted YM022 to Phase I clinical studies. Greater aqueous solubility and oral bioavailability was achieved by the introduction of basic groups, by replacing either the 5-phenyl group by a 2-pyridyl substituent, or the 3-methyl group of the aryl urea moiety by a methylamino group as in YF476¹⁶⁸ (31, also known as YM-220). This compound has shown similar binding potency at CCK_2 receptors to YM022, but 5-fold higher selectivity with respect to CCK_1 . By i.v. administration, YF476 displayed similar behavior to YM022 and was 15- and 4-fold more potent than famotidine in the inhibition of gastrin-stimulated acid secretion in anaesthetized rats and Heidenheid pouch dogs, respectively.¹⁶⁹ In this last animal model, YF476 was almost as effective by oral as by i.v. administration.¹⁶⁸ In rats, YF476 completely reversed the hypergastrinemia and cell proliferation caused by omeprazole and gastrin in ulcerated gastric mucosa.¹⁷⁰ Moreover, this antagonist showed a prolonged duration of action upon oral¹⁶⁸ and subcutaneous¹⁷¹ administration. On account of the good pharmacological profile of YF476 in *in vivo* animal models, Yamanouchi and Ferring chose to evaluate this compound for the treatment of gastro-oesophageal reflux disease (GORD), and as YM022 is in Phase I clinical studies.





Figure 4.

In an attempt to alleviate the relapse problem, frequently encountered in the chemotherapy of peptic ulcers with histamine H_2 receptor antagonists, researchers at Shionogi designed and synthesized hybrid compounds of the H_2 receptor antagonists famotidine or roxatidine and the CCK_2 antagonist L-365,260, using different spacers to joint both types of antagonists.^{172,173} Applying this strategy, they obtained dual CCK_2/H_2 antagonists such as compound 32 (Fig. 4), which showed a pA_2 value of 6.8 for the histamine H_2 receptor and an IC_{50} value of 19 nM for the CCK_2 receptor. However, although these compounds showed good gastric acid antisecretory activities by i.v. administration, in the oral route these activities were much lower, indicating low oral bioavailability.¹⁷⁴

3. Dual CCK_1/CCK_2 Receptor Antagonists

Researchers at Fujisawa postulated that dual antagonists of CCK_1 and CCK_2 receptors might be more efficacious for the treatment of pancreatitis than selective CCK_1 antagonists, based on the following reasons. CCK_1 receptor antagonists inhibit pancreatic exocrine secretion on the one hand, whilst they also stimulate gastric acid secretion, through the inhibition of the CCK_1 receptor mediated somatostatin release from D cells of the gastric mucosa, which in its turn stimulates pancreatic exocrine secretion. Additionally, it is known that lowering of pH in the duodenum by gastric acid is one of the important factors in accelerating pancreatic exocrine secretion, which is considered to be an exacerbating factor of pancreatitis. Accordingly, gastric acid secretion inhibitors such as histamine H_2 blockers and proton pump inhibitors are often prescribed for the treatment of pancreatitis. This hypothesis was supported by the fact that the joint administration of the CCK_1 antagonist FK-480 and the CCK_2 antagonist YM022 inhibited more profoundly the casein-stimulated pancreatic exocrine secretion than both compounds in separated treatment.¹⁷⁵ To study this hypothesis, several 1,4-benzodiazepine-based dual CCK_1/CCK_2 receptor antagonists were designed combining structural elements of FK-480 (20) and FR175985 (Table VI, compound 33), potent and selective antagonists of CCK_1 and CCK_2 receptors, respectively. In this design, it was considered that the incorporation of additional groups into the 9 position of the 1,4-benzodiazepine skeleton of FR175985 (33) would introduce steric repulsion between substituents at 1 and 9 positions, forcing to the 1,4-benzodiazepine system to adopt a similar structure to that in FK-480, leading to a possible dual antagonism. Among the different alkyl groups incorporated into position 9, the methyl group was chosen as it gave the best binding results at CCK_1 and CCK_2 receptors.¹⁷⁶ The first compounds of this series bound to both receptor subtypes with nanomolar potency, but had poor water solubility. To improve this issue, maintaining binding potencies, similar strategies to those previously commented for other 1,4-benzodiazepine derivatives were applied, such as introduction of acidic groups into the aryl urea moiety, as in the tetrazole derivative FR193108 (34).¹⁷⁵ This compound showed potent binding affinity at both receptors, but it was found to be only poorly absorbed upon oral administration in rats. In order to improve bioavailability, this lead compound was optimized by synthesizing analogues with a lower molecular weight, reducing the size of the substituents at positions 1 and 5. The

Table VI. Most Significant 1,4-Benzodiazepine-Derived Dual CCK₁/CCK₂ Receptor Antagonists

Compound	Config. (*)	R ¹	R ²	R ³	IC ₅₀ (nM)		Selectivity CCK ₁ /CCK ₂
					CCK ₁ ^a	CCK ₂ ^b	
FK-480 (20)	<i>S</i>	--	--	--	0.67	310	0.002
FR175985 (33)	<i>R</i>	--	--	--	62	0.087	712
FR193108 (34)	<i>R</i>				9.2	0.38	24
FR196979 (35)	<i>RS</i>		CH ₃	CH ₃	2.2	0.68	3.2
FR202893 (36)	<i>RS</i>		CH ₂ -CH ₃	CH ₃	0.9	1.6	0.56
FR208418 (37)	<i>S</i>		CH ₂ -CH ₃	CH ₃	8.2	7.8	1.05
FR208419 (38)	<i>R</i>		CH ₂ -CH ₃	CH ₃	0.3	1.0	0.3

^aInhibition of [¹²⁵I]CCC-8 binding to rat pancreatic membranes.^bInhibition of [¹²⁵I]CCC-8 binding to guinea pig cerebral cortical membranes.


5-substituent can be exchange by a methyl or ethyl group without loss in the binding affinities at both receptors. However, the size of the 1-substituent cannot be significantly reduced without decreasing the binding potency at CCK₂ receptors. The comparison of the binding data for the racemic compound **36** and both resolved enantiomers **37** and **38** shows the importance of the (*R*)-configuration at C-3 for the binding to both receptor subtypes. The activity of FR208419 (**38**) after oral administration, estimated from the ID₅₀ value (0.23 mg/kg) obtained in preliminary evaluation in gastric emptying effects, was considered to be high enough for further biological evaluations with view to clinical development.¹⁷⁷

C. Benzazepine-Based CCK₂ Antagonists

By analogy with the 1,4-benzodiazepines, 1-benzazepin-2-ones have also been used as templates in the design of CCK₂ receptor antagonists. Initial examples lacked the potency and selectivity afforded by their 1,4-benzodiazepine-based analogues,¹⁷⁸ but after a parallel structural optimization some very

potent antagonists were developed.¹⁷⁹ Thus, CP212,454 (**39**) (Table VII) exhibited higher CCK₂ receptor affinity than L-365,260 and was 367-fold selective over CCK₁ receptors. In guinea pigs, this compound potently inhibited pentagastrin-induced gastric acid secretion with a lower ED₅₀ (0.8 mg/kg s.c.) than L-365,260 (ED₅₀ = 1.5 mg/kg s.c.).¹⁷⁹ However, its low water solubility suggests a probable poor oral bioavailability. Modification of the structure of **39**, by replacing the 5-phenyl group by a cyclohexyl and inserting ionizable groups such as a carboxylic acid led to the analogue CP310,713 (**40**). This compound showed improved water solubility and *in vivo* efficacy (ED₅₀ = 0.03 mg/kg s.c. in the pentagastrin-induced gastric acid secretion model).¹⁸⁰ However, its low efficacy in preclinical studies led to stop the development of CP310,713 (**40**).⁸³

Table VII. Significant 1-Benzazepine-2-one-Based CCK₂ Receptor Antagonists




Compound	R ¹	R ²	IC ₅₀ (nM)		Selectivity CCK ₁ /CCK ₂	Reference
			CCK ₁ ^a	CCK ₂ ^b		
CP212,454 (39)		Cl	176	0.48	367	179
CP310,713 (40)		CO ₂ H	1,400	0.10	14,000	180

^aInhibition of [¹²⁵I]CCC-8 binding to guinea pig pancreatic membranes.

^bInhibition of [¹²⁵I]CCC-8 binding to guinea pig cortex.

D. 1,5-Benzodiazepine-Based CCK₂ Antagonists

The introduction of functional assays, along with radioligand binding assays, in high throughput screenings by Glaxo's researchers allowed them the discovery of a series of 1,5-benzodiazepine derivatives as the first non-peptide CCK₁ receptor agonists.¹⁸¹ In these random screenings, some of the compounds showed CCK₂ antagonist activity, which was optimized introducing substituents into the 1,5-benzodiazepine skeleton similar to those found successful in the Merck 1,4-benzodiazepine antagonists. Lead compounds were obtained by positioning an aryl urea moiety at C-3 and locating branched or bulky groups at N-1 and N-5 positions. Structure-activity relationship (SAR) studies, as in the 1,4-benzodiazepine series, showed a similar influence of the absolute configuration at C-3 on the binding potency, being the (*R*)-enantiomers more potent at CCK₂ receptors.¹⁸² Among these compounds GV150013X (Table VIII, compound **41**) was the first selected for pharmacological development. In the guinea pig myenterium plexus, where both CCK₂ and CCK₁ receptors are found GV150013X antagonized the CCK-4-induced contractions with a pK_B of 8.9, while for the inhibition of CCK₁ agonist-induced contractions the pK_B was 5.9. In the isolated gastric mucosa, this compound also showed antagonistic activity in the pentagastrin-induced gastric acid secretion, although was less potent with a pK_B of 7.4.¹⁸² *In vivo*, GV150013X has shown dose-related anxiolytic effects in several animal models.¹⁸³ In all the anxiety models used, this antagonist displayed similar efficacy to the dipeptoid CCK₂ antagonist PD134,308 and to the standard benzodiazepine diazepam, without tolerance or rebound anxiogenesis upon withdrawal after chronic treatment (7 days at 0.3 µg/kg, p.o.) observed with diazepam. In general pharmacological studies, GV150013X did not show effect up to 3 mg/kg, p.o. in the rota-rod test, in the passive avoidance, and on pentobarbitone sleeping time.¹⁸²

Table VIII. More Significant 1,5-Benzodiazepine-Based CCK₂ Receptor Antagonists

Compound	Config. (*)	R ¹	R ²	R ³	Affinity		Selectivity CCK ₁ /CCK ₂	Ref.
					CCK ₁	CCK ₂		
GV150013X (41)	R		Phenyl	Phenyl	pK _i ^a 6.15	pK _i ^b 8.64	309	182
GV191869X (42)	R			Phenyl	pK _i ^a 5.70	pK _i ^b 9.40	4,988	81
GV199114X (43)	S		Cyclohexyl		pK _i ^a 7.10	pK _i ^b 8.60	40	185
44	--				IC ₅₀ ^c 500	IC ₅₀ ^d 2.0	250	186
45	--				IC ₅₀ ^c 460	IC ₅₀ ^d 6.0	77	186

^apK_i values of inhibition of the binding of [³H]CCK-8 to rat pancreatic membranes.

^bpK_i values of inhibition of the binding of [³H]CCK-8 to guinea pig cerebral cortex membranes.

^cIC₅₀ nM values of the binding of [propionyl-³H]CCK-8 to mouse pancreas.

^dIC₅₀ nM values of the binding of [propionyl-³H]CCK-8 to mouse cortical membranes.

Additionally, GV150013X improved sleep in aged rats, without detecting tolerance after chronic treatment, while development of tolerance towards benzodiazepines was monitored following chronic treatment with triazolam.¹⁸⁴ Based on these results, GV150013X has progressed to Phase II clinical trials for anxiety and sleep disorders, although no data are currently available.







In order to improve physicochemical properties of GV150013X, the phenyl group at the N-5 position was replaced by a morpholinoethyl substituent to give GV191869X (42). This analogue showed improved solubility and was a more potent and selective CCK₂ antagonist.⁸¹ GV191869X (42) also exhibited anxiolytic activity in both the mouse light/dark box and in the marmoset 'human threat' test models, maintaining a significant effect in the dose range of 0.01–10 µg/kg.⁸¹ Similarly, GV199114X (43) also showed improved aqueous solubility and oral bioavailability. This antagonist was equipotent by i.v. (60% at 0.3 mg/kg) and oral administration in the inhibition of pentagastrin-induced gastric acid secretion in rats.¹⁸⁵ Although based on the *in vivo* potency and plasma pharmacokinetics this compound could be a suitable candidate to investigate the actions of peripheral CCK₂ receptors, this may be hampered by the low receptor selectivity margin (40-fold, based on radioligand binding affinities).

By choosing identical substituents at both N-1 and N-5 positions of the 1,5-benzodiazepine template, Shionogi avoided the need for enantiomer resolution or stereoselective synthesis. Optimization of this substituent identified cyclopropylcarbonyl-methyl as the preferred one. When this substituent was combined with the introduction of acid groups into the aryl urea moiety, such as the thioacetic acid of compound 44, the CCK₂ receptor affinity was comparable to that of L-365,260.¹⁸⁶ However, *in vivo*, the optimum compound of this series was the ethyl ester derivative 45 which showed potent inhibition of pentagastrin-induced gastric acid secretion in anaesthetized rats, with an ED₅₀ value of 0.06 mg/kg upon intraduodenal administration, being almost twice more potent than L-365,260 and YM022 in this assay.¹⁸⁶

E. Ureidoacetamide CCK₂ Antagonists

The ureidoacetamide-based CCK₂ receptor antagonists may be considered acyclic analogues of the 1-carbonylmethyl-1,4-benzodiazepine-derived antagonists, resulting from the opening of the C₃-N₄ bond. Most of these ureidoacetamide analogues exhibited similar biological properties to those of their 1,4-benzodiazepine parent compounds. The first of these compounds were described by Rhône-Poulenc,¹⁸⁷ such as RP 69758 (Table IX, compound 46), which displayed comparable binding potency in guinea pig cortex to L-365,260, but based on ex vivo binding studies did not significantly penetrate the CNS.¹⁸⁸ The replacement of the acetic acid moiety in R² by ethyl sulphonate and the introduction of a methoxy group into the phenyl group led to a 10-fold increase in the CCK₂ receptor affinity and selectivity for RP 73870 (47).¹⁸⁹ This increase in the *in vitro* affinity was also observed in the *in vivo* assays. This compound was 10-fold more potent than the dipeptoid CI-988 in the inhibition of gastrin-stimulated gastric acid secretion in the in situ perfused rat stomach

Table IX. Significant Ureidoacetamido-Based CCK₂ Receptor Antagonists

Compound	R ¹	Ar	R ²	Affinity		Selectivity CCK ₁ /CCK ₂	Reference
				CCK ₁	CCK ₂		
RP 69758 (46)	NHPh	Ph	CH ₂ CO ₂ H	K _i ^a 1,254	K _i ^b 9.0	139	188
RP 73870 (47)	NMePh			K _i ^a 1,634	K _i ^b 0.48	3,404	189
RPR1011367 (48)			CH ₂ CO ₂ H	--	K _i ^b 3	--	83
S-0509 (49)	OtBu		CH ₂ CO ₂ Na	IC ₅₀ ^c 3,400	IC ₅₀ ^d 42	81	190
DA-3934 (50)	NMePh		CH ₂ CO ₂ H	IC ₅₀ ^e 877	IC ₅₀ ^f 0.4	2,193	192
D51-9927 (51)			CH ₂ CO ₂ H	IC ₅₀ ^e 172	IC ₅₀ ^f 0.06	2,867	193

^aK_i (nM) inhibition of the binding of [propionyl-³H]CCK-8 to guinea pig pancreas.

^bK_i (nM) inhibition of the binding of [propionyl-³H]CCK-8 to guinea pig cortex.

^cIC₅₀ (nM) inhibition of the binding of [propionyl-³H]CCK-8 to mouse pancreas.

^dIC₅₀ (nM) inhibition of the binding of [propionyl-³H]CCK-8 to mouse cortical membranes.

^eIC₅₀ (nM) inhibition of the binding of [¹²⁵I]CCK-8 to human CCK₁ receptors.

^fIC₅₀ (nM) inhibition of the binding of [¹²⁵I]gastrin to human gastrin receptors.

(ID₅₀ = 0.05 mg/kg i.v., 3 mg/kg p.o.). Unlike RP 69758, compound **47** did not inhibit basal acid secretion at concentrations required to inhibit the gastrin-induced secretion. Moreover, in models of gastric ulceration in rats, RP 73870 specifically inhibited acid-dependent mechanisms, being as effective as the proton pump inhibitor omeprazole and ineffective in models of ethanol-, restraint stress- or insulin-induced ulceration. Rhône-Poulenc also designed restricted analogues of their ureidoacetamides such as RPR1011367 (**48**) to investigate CCK₂-mediated behavioural effects in animal models. This antagonist inhibited the CCK-8-stimulated firing of rat hippocampal neurons with a 4- and 10-fold higher potency than CI-988 and L-365,260, respectively, and this higher potency was also observed *in vivo*, where this compound displayed anxiolytic-like effects in the elevated X-maze test.⁸³


Shionogi also designed ureidoacetamides in which a benzophenone moiety was introduced to keep a greater similarity with the 1,4-benzodiazepine structure.¹⁹⁰ Among the compounds of this series, S-0509 (**49**) afforded the optimum balance of receptor affinity and selectivity and *in vivo* potency. On anaesthetized pylorus-ligated rats this antagonist was about 30- and 200-fold more potent than L-365,260 by i.v. (ED₅₀ = 0.001 mg/kg) and intraduodenal (0.003 mg/kg) administration, respectively. Unlike L-365,260 and YM022, S-0509 did not enhance morphine analgesia on the tail flick test. This result prompted Shionogi researchers to conclude that S-0509 has poor blood-brain permeability and, because of that, could be a good tool to distinguish between peripheral and central effects of CCK₂ antagonism. S-0509 is currently found at Phase I clinical trials for gastric secretion disorders.

Researchers at Daiichi have also reported on ureidoacetamide-based CCK₂ antagonists that include a phenoxyacetic acid moiety in their structure.¹⁹¹⁻¹⁹³ For example, DA-3934¹⁹² (**50**) and D51-9927¹⁹³ (**51**), which potently inhibited the [¹²⁵I]gastrin binding to CHO cells transfected with human CCK₂ receptors. In lumen-perfused anaesthetized rats D51-9927 inhibited pentagastrin-stimulated acid secretion by i.v. (ED₅₀ = 0.004 mg/kg) and intraduodenal routes (ED₅₀ = 0.5 mg/kg). Chronic administration (28 days) of this antagonist to rats (3, 30 mg/kg/day, p.o.), together with omeprazole (60 mg/kg/day, p.o.), inhibited the hypersecretory response to gastrin-G17, otherwise observed on cessation of omeprazole treatment.⁸³

F. Quinazolinone-Based CCK₂ Antagonists

The strategy of manipulating the asperlicin structure, by appropriate bond disconnections, retaining the quinazolinone and indole templates, led Lilly's researchers to the discovery of another family of potent and selective CCK₂ antagonists.¹⁹⁴ The lack of chiral centers in this series of compounds significantly facilitated the structure optimization, from which LY-202769 (Table X, compound **52**) was selected as the best compound, although the reported pharmacological results of this compound are limited to its effect on midbrain dopamine activity.¹⁹⁵ The reported SAR study showed the importance of the isopropoxy substituent at the *meta* position of the 3-phenyl group as well as the size and flexibility of the carbon chain linker between the quinazolinone and indole templates.^{194,196}

Parke-Davis also pursued this design approach, maintaining the quinazolinone skeleton, but replacing the 3-alkyl-indole moiety by an aryl urea. The first compounds of this series, such as **53** and **54**, showed only moderate binding potency and selectivity at CCK₂ receptors,¹⁹⁷ which were significantly improved by replacing the methylene bridge by a NH, as in compounds **55** and **56**.¹⁹⁸ As in LY-202769, the presence of the *meta*-isopropoxy group, or other of similar size as the dimethylamino of **56**, is important both for the potency and selectivity. The presence of electron-withdrawing substituents in position 3 of the phenylurea group also significantly enhanced the CCK₂ potency. Both compounds **55** and **56** showed anxiolytic-like effects in the rat X-maze test after oral administration, although their lower effective dose (1.0 mg/kg) was 10-fold higher than that of the reference compound LY-202769 (**52**) (0.1 mg). Both compounds **55** and **56** showed poor aqueous solubility and, as consequence, the estimated bioavailability of **55** was < 5%, while that of **56** was 22% after oral administration.¹⁹⁸

Table X. Significant Quinazolinone-Based CCK₂ Receptor Antagonists

Compound	X	R ¹	R ²	IC ₅₀ (nM)		Selectivity CCK ₁ /CCK ₂	Reference
				CCK ₁ ^a	CCK ₂ ^b		
LY-207269 (52)	--	--	--	>10 ⁴	9.3	>1,075	194
53	CH ₂	CH ₃	OPr	1,637	879	1.9	197
54	CH ₂	CO ₂ Et	OPr	1,465	126	11.6	197
55	NH	CO ₂ tBu	OPr	2,960	1.9	1557	198
56	NH	CN	N(Me) ₂	7,100	14	507	198


^aIC₅₀ for inhibition of [³H]-L-364,718 (**52**) or [¹²⁵I]CCK-8 binding to rat pancreas.

^bIC₅₀ for inhibition of [¹²⁵I]CCK-8 binding to mouse brain membranes.

G. Dipeptoids

Researchers at Parke-Davis used the C-terminal tetrapeptide sequence of CCK (CCK-4) as a starting point for the rational design of CCK₂ receptor ligands. This design was based on the finding that the Trp and Phe residues were the minimum and necessary sequence to impart micromolar affinity at CCK₂ receptors¹⁹⁹ (Table XI). They adopted the strategy of sequentially optimizing both the N- and C-terminal residues of the Boc-Trp-Phe-NH₂ dipeptide. The resulting dipeptide analogues were designed as dipeptoids.²⁰⁰ As shown in Table XI, the N-terminus SAR studies revealed that the replacement of Trp by α -methyl-tryptophan, together with bulky cycloalkyl carbamate groups at this residue were preferred. Among these bulky groups, the 2-adamantyloxycarbonyl was the optimum.²⁰⁰ These studies also showed that modifications at the tryptophan indole ring were not well tolerated.²⁰¹ Keeping constant in the structure the N-(2-adamantyloxycarbonyl)- α -methyltryptophan residue, the C-terminus optimization was focused on the incorporation of carboxylic acid-containing groups at positions 1 and 2 of the 2-phenylethylamine moiety.²⁰² CI-988 (also named PD-134,308, **58**) emerged from this SAR study as the most potent and selective CCK₂ antagonist of this series. This compound and its analogue CAM1189 (**59**) blocked the pentagastrin-stimulated gastric acid secretion by iv infusion of 0.5 and 0.07 μ mol/kg, respectively.²⁰² Compound **59** was equipotent by i.v. and s.c. administration (ED₅₀ = 0.07 μ mol/kg), being more potent than the H₂ receptor antagonist ranitidine (ED₅₀ = 0.19 μ mol/kg). The replacement of the CI-988 carboxylic acid by some bioisosteric groups, such as the sulphonic acid of compound **60** did not significantly affected the CCK₂ binding affinity, although these replacements reduced the CCK₁/CCK₂ selectivity of the resulting compounds.²⁰³ Based on its higher CCK₂ selectivity versus CCK₁ receptors, CI-988 was chosen for studies on its behavioral effects on the CNS. In these studies, CI-988 exhibited potent anxiolytic effects in several animal models of anxiety, including the mouse black/white box test, the rat plus maze and social interaction tests, and the marmoset human threat test.²⁰⁴ These effects were comparable in magnitude to those produced by diazepam, but, unlike diazepam, CI-988 did not produced sedation, and there was no evidence of development of tolerance or any sign of withdrawal anxiogenesis after abrupt termination of the treatment. However, these anxiolytic effects have not been confirmed in Phase II clinical trials in patients with generalized anxiety and panic disorders.^{205–208} Neither has CI-988

Table XI. Significant Dipeptoid CCK Receptor Antagonists



Compound	Config.			R ¹	R ²	R ³	R ⁴	IC ₅₀ (nM)		Selectiv ^c	Ref.
	*	1	2					CCK ₁ ^a	CCK ₂ ^b		
CCK-8								0.12	0.27	0.44	
CCK-4								5,330	2.6	2,050	
BocTrp-PheNH ₂								--	73,000 ^d	--	199
57	RS	--	--	1-Ad ^e	H	H	H	--	5,000 ^d	--	200
CI-988 (58) (PD-134,308)	R	--	R	2-Ad ^f	H	HNCO(CH ₂) ₂ CO ₂ H	H	4,300	1.7	2,529	202
CAM1189 (59) (PD-136,450)	R	--	R	2-Ad	H	HNCOCH=CHCO ₂ H	H	709	0.7	1,100	202
60	R	--	R	2-Ad	H	HNCO(CH ₂) ₂ SO ₃ Na	H	1,010	1.3	780	203
PD-135,666 (61)	R	S	--	2-Ad	CH ₂ CO ₂ H	H	H	25.1	0.15	170	219
CAM 1481 (62) (PD-140,548)	S	R	--	2-Ad	CH ₂ CO ₂ H	H	H	2.82	260	0.01	219
PD-140,547 (63)	S	S	--	2-Ad	CH ₂ CO ₂ H	H	H	539	13.2	41	219
PD-140,723 (64)	R	R	--	2-Ad	CH ₂ CO ₂ H	H	H	186	9.3	20	219
PD-149,164 (65)	R	S	--	2-Ad	CH ₂ CO ₂ H	H	F	75	0.08	938	220
66	R	S	--	2-Ad	CH ₂ CO ₂ H	H	NO ₂	225	0.19	1,184	220
67	S	R	--		CH ₂ CO ₂ H	H	H	7.9	1,160	0.007	219
68	R	S	--		CH ₂ CO ₂ H	H	H	3.9	4.2	1	219
CI-1015 (69)	R	S	S	2-Ad	--	--	--	2,900	3.0	967	215

^aIC₅₀ for inhibition of [¹²⁵I]CCK-8 binding to rat pancreas membranes.^bIC₅₀ for inhibition of [¹²⁵I]CCK-8 binding to mouse cortex membranes.^cSelectivity, CCK₁/CCK₂.^dK_i value for the inhibition of [³H]-Boc-β-alanyl-CCK-4 binding to mouse cortex membranes.^e1-Ad = 1-adamantyl.^f2-Ad = 2-adamantyl.







shown significant anxiolytic effects on lactate-²⁰⁹ and CCK-4-induced²¹⁰ panic symptoms in healthy volunteers. These poor results, along with the reported CCK₂ partial agonist activity on histidine decarboxylase in rat stomach enterochromaffin-like cells²¹¹ and on stomach histamine release and acid secretion,²¹² as well as the full CCK₁ agonist activity on the amylase release in the rat pancreas,²¹³ have hampered the further clinical development of CI-988. In diabetic rats, CI-988 enhanced the analgesia induced by morphine,²¹⁴ and induced antinociceptive effects on mechanical hyperalgesia.²¹⁵

The lack of anxiolytic effects of CI-988 in humans has been attributed to its low bioavailability,^{210,216} in part due to its high molecular weight. Therefore, to improve the pharmacokinetic profile, the structure of CI-988 was modified with the aim of reducing the molecular weight and improving the aqueous solubility and absorption, maintaining the binding potency and selectivity at CCK₂ receptors. As the *N*-(2-adamantylloxycarbonyl)- α -methyl-D-tryptophan moiety is critical for affinity, modifications were focused on the C-terminus.²¹⁶ These modifications led to the identification of the analogue CI-1015 (Table XI, 69),²¹⁶ which showed similar CCK₂ binding affinity that CI-988, although with lower selectivity versus CCK₁ receptors. This compound exhibited CCK₂ antagonist profile in the rat ventromedial hypothalamus assay with a K_e of 34 nM. It also showed an anxiolytic like profile orally in the standard anxiety (X-maze) paradigm with a minimum effective dose of 0.1 μ g/kg. Although CI-1015 was less water soluble than CI-988, the oral bioavailability in rats was improved nearly 10 times when dosed as a solution in hydroxypropyl- β -cyclodextrin. The blood-brain permeability was also enhanced relative to CI-988. On the basis of the overall improved pharmacokinetic profile,¹¹⁷ CI-1015 was chosen as a development candidate, although results on clinical studies have not been reported.

Additional SAR studies have demonstrated the importance of the central amide bond between the N- and C-terminus of these dipeptoids for their CCK₂ binding potency and selectivity.¹¹⁸ On the other hand, the diastereoisomeric dipeptoids substituted at position 1 of the phenyl-ethylamine moiety **61–64** (Table XI) have proved the stereochemical preference for each receptor subtype.²¹⁹ Thus, the stereoisomer **61** (PD-135,666) was a potent and selective CCK₂ antagonist in the electrophysiological test on the rat ventromedial nucleus of the hypothalamus, with a K_e value of 2.8 nM, and was also anxiolytic in the mouse light/dark box test with a minimum dose of 0.01 mg/kg, s.c. On the contrary, its enantiomer **62** (PD-140,548) was a selective and competitive CCK₁ antagonist, which inhibited the CCK-8-evoked amylase release from rat pancreatic acinar cells with a K_e value of 16 nM.²¹⁹ In the *trans*-2-methylcyclohexyl derivatives **67** and **68** the reversal of the stereochemistry transformed the CCK₂ antagonist **67** into the mixed CCK₁/CCK₂ antagonist **68**, which showed antagonist properties in both CCK₁ and CCK₂ models. Introduction of small electron withdrawing substituents, such as F or NO₂, into the *para* position of the phenyl group of compound **61** increased affinity and selectivity at CCK₂ receptors in the analogues **65** and **66**.²²⁰ It is interesting to note that compound **65** (PD-149,164) was a potent antagonist at CCK₂ receptors and showed full CCK₁ agonist activity *in vivo* in the exocrine pancreas of rats and *in vitro* in rat pancreatic acini, while its enantiomer was a CCK₁ antagonist.²²¹

With the aim of developing pharmacophore models of the dipeptoid-CCK receptor recognition or proving conformational hypothesis, conformationally constrained analogues have also been designed. Among these analogues, the use of a proline ring to constrain the tryptophan residue led to a significant decrease in the binding potency at CCK₂ receptors and reversal of the selectivity towards CCK₁ receptors in compound **70** (Table XII).²²² Perhaps the low affinities of this compound could have been improved after the resolution of the epimeric mixture at the proline stereogenic center. Restriction of the phenyl group rotation in CI-988, by incorporation of a tetrahydronaphthyl group into the phenyl-ethylamine moiety, led to a decrease of approximately 10-fold in the CCK₂ selectivity for the diastereoisomeric analogues **71** and **72**.²²³ The proline ring has also been used to restrict the C-terminus in compounds **73–75**.²²⁴ These compounds also displayed reduced CCK₂ affinity and selectivity, but it is worthy of note that, although the three compounds exhibited comparable CCK₂

Table XII. Conformationally Constrained Dipeptoid CCK Receptor Antagonists

Compound	Afinity (nM)		Selectivity CCK ₁ /CCK ₂	Reference																										
	CCK ₁	CCK ₂																												
<div></div> <div>70</div> <div>(2-Adoc)</div>	1,050 ^a	2,080 ^b	0.50	222																										
<div></div> <div>71</div> <div>(2-Adoc)</div>	<div>Configuration</div> <table><tr><th>N°</th><th>1</th><th>2</th></tr><tr><td>71</td><td>S</td><td>R</td></tr><tr><td>72</td><td>R</td><td>R</td></tr></table>		N°	1	2	71	S	R	72	R	R	460 ^a	2.31 ^b	199	223															
N°	1	2																												
71	S	R																												
72	R	R																												
<div></div> <div>73</div> <div>(2-Adoc)</div>	<table><tr><th>N°</th><th>R¹</th><th>R²</th></tr><tr><td>73</td><td>Cl</td><td>Cl</td></tr><tr><td>74</td><td>F</td><td>F</td></tr><tr><td>75</td><td>H</td><td>NO₂</td></tr></table>		N°	R ¹	R ²	73	Cl	Cl	74	F	F	75	H	NO ₂	1,690 ^a	6.16 ^b	274	223												
N°	R ¹	R ²																												
73	Cl	Cl																												
74	F	F																												
75	H	NO ₂																												
<div></div> <div>76</div> <div>(2-Adoc)</div>	<table><tr><th>N°</th><th>*</th></tr><tr><td>76</td><td>S</td></tr><tr><td>77</td><td>R</td></tr></table>		N°	*	76	S	77	R	749 ^c	45.7 ^d	16.4	224																		
N°	*																													
76	S																													
77	R																													
			394 ^c	17.6 ^d	22.4	224																								
			2,104 ^c	26.3 ^d	80	224																								
<div></div> <div>78</div> <div>(2-Adoc)</div>	<div>Configuration</div> <table><tr><th>N°</th><th>2</th><th>S</th><th>11b</th><th>*</th><th>n</th></tr><tr><td>78</td><td>S</td><td>S</td><td>R</td><td>S</td><td>0</td></tr><tr><td>79</td><td>R</td><td>R</td><td>S</td><td>S</td><td>0</td></tr><tr><td>80</td><td>S</td><td>S</td><td>R</td><td>R</td><td>1</td></tr></table>		N°	2	S	11b	*	n	78	S	S	R	S	0	79	R	R	S	S	0	80	S	S	R	R	1	18 ^a	0.30 ^b	60	225
N°	2	S	11b	*	n																									
78	S	S	R	S	0																									
79	R	R	S	S	0																									
80	S	S	R	R	1																									
						3.9 ^a	60 ^b	0.065	225																					
<div></div> <div>78</div> <div>(2-Adoc)</div>	<table><tr><th>N°</th><th>2</th><th>S</th><th>11b</th><th>*</th><th>n</th></tr><tr><td>78</td><td>S</td><td>S</td><td>R</td><td>S</td><td>0</td></tr><tr><td>79</td><td>R</td><td>R</td><td>S</td><td>S</td><td>0</td></tr><tr><td>80</td><td>S</td><td>S</td><td>R</td><td>R</td><td>1</td></tr></table>		N°	2	S	11b	*	n	78	S	S	R	S	0	79	R	R	S	S	0	80	S	S	R	R	1	4.7 ^e	>10,000 ^f	<0.0005	227
N°	2	S	11b	*	n																									
78	S	S	R	S	0																									
79	R	R	S	S	0																									
80	S	S	R	R	1																									
						1.7 ^e	202 ^f	0.008	227																					
						7.4 ^e	2,700 ^f	0.003	226																					

^aIC₅₀ for inhibition of the [¹²⁵I]CCK-8 binding to rat pancreas membranes.^bIC₅₀ for inhibition of the [¹²⁵I]CCK-8 binding to mouse cortex membranes.^cK_i values for inhibition of the [propionyl-³H]CCK-8 binding to guinea pig pancreas membranes.^dK_i values for inhibition of the [propionyl-³H]CCK-8 binding to guinea pig cortex membranes.^eIC₅₀ for inhibition of the [propionyl-³H]CCK-8 binding to rat pancreas membranes.^fIC₅₀ for inhibition of the [propionyl-³H]CCK-8 binding to rat cortex membranes.


affinities on guinea pig cortex and on rat brain CCK₂ receptor expressed in CHO cells, they inhibited the CCK-8-induced inositol phosphate production in these cells with rather different IC₅₀ values (37.4, 389, and 507 nM for **73–75**). The approximate 10-fold discrepancy observed in the antagonistic activity of compound **73** with respect to **74** and **75**, has been attributed to differences in the binding at two affinity states of CCK₂ receptors.²²⁴ As shown in Table XII, the enantiomeric compounds **76** and **77**, where a α,β -didehydrotryptophan replaces the α -methyl-tryptophan, provided additional evidence of the stereoselectivity in the binding to both receptor subtypes.²²⁵ The highly CCK₁ selective compounds **78–80** were designed based on the hypothesis that the folding of the peptide backbone of dipeptoids into a β -turn-like conformation could contribute to their bioactive conformation at CCK₁ receptors.^{226,227} These compounds incorporate the 2-amino-3-oxohexahydroindolino[8,7-*b*]indole-5-carboxylate scaffold, a probed β -turn mimetic, as replacement of the α -methyl-tryptophan. This design strategy has provided the most potent and selective CCK₁ antagonists in the dipeptoid series.

H. Dibenzobicyclo[2.2.2]octane and Bicycloheteroaromatic Scaffold-Based CCK₂ Antagonists

James Black Foundation' researchers have also designed a series of selective CCK₂ receptor antagonists based on the topography of the CCK-4 structure. They looked for rigid skeletons that

could replace the peptide backbone of this tetrapeptide while maintaining the stereoelectronic features obtained from molecular mechanic calculations and fluorescence studies on CCK-4. From this basis, they proposed that the two aromatic rings of the tryptophan and phenylalanine side chains of CCK-4 are interacting in a π -stacking arrangement with a separation of 5–7 Å, and designed the dibenzobicyclo[2.2.2]octane (BCO) scaffold as mimic of the peptide backbone.²²⁸ One of the fused phenyl rings of this skeleton could mimic one of the aromatic side chains, and a second aromatic group would be appended as part of the R¹ moiety in compounds **81–85** (Table XIII). Some of the initial compounds of this series, exemplified by the proline derivatives **81** and **82**, exhibited submicromolar affinity for CCK₂ receptors, and selectivity > 60-fold over the CCK₁ receptors.²²⁸ The first SAR

Table XIII. Dibenzobicyclo[2.2.2]octane and Bicycloheteroaromatic Scaffold-Based CCK₂ Receptor Antagonists



Compound	X	Y	R ¹	R ²	pK_i		Selectiv. CCK ₁ /CCK ₂	Ref.
					CCK ₁ ^a	CCK ₂ ^b		
81	--	--	1-Ad ^c	D-Pro-Gly	4.83	6.67	69	228
82	--	--	1-Ad ^c	L-Pro-D-Ala	4.62	7.39	589	228
83	--	--	1-Ad ^c		5.68	8.80	1,318	229
84	--	--	Cyclohexyl		5.30	7.76	288	229
85	--	--	2-Naphtyl		5.79	8.57	603	229
JB93182 (86)	NH	CH ₂	H	CO ₂ H	5.44	8.96	3,311	230
87	S	CH ₂	H	CO ₂ H	5.65	8.91	1,820	230
88	NH	N	H	CO ₂ H	5.83	8.28	282	230
89	NH	CH ₂	F	CO ₂ H	5.48	9.13	4,467	230
90	NH	CH ₂	H		5.89	8.92	1,072	230
91	NH	N	--	CO ₂ H	4.74	8.30	3,631	234

^a pK_i values for the inhibition of the binding of [¹²⁵I]CCK-8 to guinea pig pancreas.

^b pK_i values for the inhibition of the binding of [¹²⁵I]CCK-8 to mouse cortical homogenates.

^cAd, adamantyl.

studies on this series showed that, among several bulky alicyclic and aromatic groups incorporated into R^1 , the 1-adamantyl group was the optimum. The replacement of the proline residue by L-phenylalanine and the glycine or alanine residues by the 3,5-dicarboxyanilide group, to give compounds **83–85**, produced an increase of, at least, one order of magnitude in the binding affinity at CCK_2 receptors. At least one of the carboxylic groups of these compounds is crucial for high binding affinity, and altering the chain length of the phenylalanine residue is detrimental to the binding.²²⁹ Interestingly, compounds **83–85** showed species-variation when evaluated *in vivo* for their ability to antagonize pentagastrin-stimulated gastric acid secretion in rats and dogs. For example, the adamantyl derivative **83**, when administered intravenously, was 700-fold less active in chronic gastric fistula dogs than in Ghosh and Schild anaesthetized rats.²³⁰ This discrepancy prompted further modification to identify a replacement for the BCO framework. This research resulted in a new series of *ortho*-disubstituted bicyclic heteroaromatic analogues which maintained the affinity and selectivity demonstrated by the BCO derivatives, but gave a more consistent *in vivo* profile.²³⁰ Thus, for example, the indole derivative JB93182 (Table XIII, compound **86**) was as potent as the BCO analogue **83** in the pentagastrin-stimulated gastric acid secretion model in anaesthetized rats, but exhibited similar potency in chronic gastric fistula dogs. The *in vivo* species-variable behavior exhibited by the BCO derivatives could be due to the interspecies variation in CCK receptors already commented in the introductory heading 2. JB93182 dose-dependently inhibited the gastrin-stimulated histidine decarboxylase activation in intact fasted rats with a ID_{50} value of 8 nmol/kg i.v., 4-fold lower than that shown by the benzodiazepine derivative YF476.²³¹ Similarly, JB93182 inhibited the gastrin-induced secretion of pancreastatin, a chromogranin A-derived peptide, from isolated rat enterochromaffin-like (ECL) cells with a IC_{50} value of 9.8 nM.²³² Both activities are related with the blockage of CCK_2 receptors in ECL cells.²³³

To study the conformational preference of the phenylalanine side chain, this residue was replaced in compound **88** by conformationally restricted phenylalanine analogues, which integrate the phenyl and amino groups into different size rings.²³⁴ In general, this structure modification was detrimental for the binding at CCK_2 receptors, except for the analogue **91** which showed similar *in vitro* profile to that of its model **88**.

Based on the same design strategy that led to the BCO-derived antagonists, a new series of antagonists based on the 2,7-dioxo-2,3,4,5,6,7-hexahydro-1-*H*-benzo[*h*][1,4]-diazonine scaffold has recently been reported.²³⁵ The initial compounds of this series, such as compounds **92** and **93** (Fig. 5), have shown moderate affinities at CCK_2 receptors, similar to that exhibited by the first members of the BCO family.

It is remarkable that a simple *N*-methylation at the indolic N–H in JB93182 (**86**) gave rise to an agonist activity of acid secretion stimulation in isolated lumen-perfused mouse stomach, without significantly affecting the binding affinity at CCK_2 receptors.²³⁶ This easy switch from antagonist to agonist functionality suggests that both the agonist and the antagonist might share a common recognition site at the receptor.

I. 1,3-Dioxoperhydropyrido[1,2-*c*]pyrimidine-Based CCK_1 Antagonists

The CCK-4 structure was also the starting point for the search of CCK antagonists at the Medicinal Chemistry Institute of Madrid. In this case, researchers focused their attention on the design of









Figure 5.

conformationally restricted analogues, using bicyclic lactam skeletons as replacements for the central dipeptide [Met³¹-Asp³²] of CCK-4. Some of the first compounds resulting from this approach, which included a 3-oxoindolizidine skeleton as a spacer between the Trp³⁰ and Phe³³ residues, represented by compound **94** (Table XIV), showed micromolar affinity at CCK₁ or CCK₂ receptors depending on the stereochemistry.²³⁷ Based on these results, the structure of the initial compounds was manipulated by replacing the 3-oxoindolizidine skeleton by analogue rigid frameworks which could impart to the aromatic side chains a more appropriate spatial orientation at the receptor recognition site. Particularly, the use of the 1,3-dioxoperhydropyrido[1,2-*c*]pyrimidine scaffold led to the discovery of a family of highly potent and selective CCK₁ receptor antagonists.²³⁸ The prototype of this group of antagonists IQM-95,333 (**95**) showed nanomolar CCK₁ affinity, comparable to that shown by devazepide (**18**) in the same assay, but with a higher selectivity, as it was devoid of affinity at brain CCK₂ receptors at the highest concentration assayed (10⁻⁵ M). This compound inhibited the CCK-8-stimulated amylase release from rat pancreatic acini with similar potency (IC₅₀ = 21.3 nM) to that of devazepide (IC₅₀ = 25.4 nM).²³⁹ As a CCK₁ antagonist, IQM-95,333 also inhibited the contractile response induced by CCK-8 in the guinea pig ileum with a comparable potency (*p*K_B = 8.4) to that of devazepide (*p*K_B = 8.5). Low doses (50–100 µg/kg, i.p.) of this antagonist, blocked the hypophagia and the hypolocomotion induced by systemic administration of CCK-8 in rats, two effects associated with the stimulation of peripheral CCK₁ receptors. Interestingly, IQM-95,333 exhibited an anxiolytic-like profile in the light/dark exploration test in mice over a wide dose range (10–5,000 µg/kg, i.p.), while the CCK₁ and CCK₂ antagonists devazepide and L-365,260, respectively, were effective only within a more limited dose range (2–100 µg/kg and 2–100 µg/kg i.p.). The three antagonists IQM-95,333, devazepide and L-365,260 also showed anxiolytic-like effects in the rat punished-drinking test (Vogel test), although within a narrow dose range.²³⁹ Although the anxiolytic effects of devazepide have previously been attributed to the blockage of CCK₂ receptors at high doses (IC₅₀ = 270 nM¹³³),²⁴⁰ this explanation can not be applied to IQM-95,333, as this compound is devoid of affinity at this receptor subtype. Therefore, these results support previous suggestions that CCK₁ receptors may also be involved in anxiogenesis.^{241–243}

To define the pharmacophore of this perhydropyrido[1,2-*c*]pyrimidine-based family of CCK₁ antagonists a thorough SAR study on the three structural domains (tryptophan, central bicyclic skeleton, and substituent at N2-position) was carried out. This study showed the importance of the (4a*S*,5*R*)-stereochemistry at the 1,3-dioxoperhydropyrido[1,2-*c*]pyrimidine template (compare compounds **95–98**, Table XIV) and the *L*-configuration at the tryptophan residue as essential requirements for CCK₁ binding affinity and subtype receptor selectivity.²³⁸ The presence of the *tert*-butoxycarbonyl group (Boc) and the amide bond, or an appropriate H-bonding surrogate such as ψ[CH(CN)NH] (**105**), were also critical for the binding at CCK₁ receptors, while the replacement of the tryptophan residue by other aromatic amino acids, such as Phe (**99**) or α-Me-Trp (**100**) led to more than 10-fold decrease in the binding potency.²⁴⁴ Interestingly, the replacement of the acid-labile Boc group by their bioisosters *tert*-butylaminocarbonyl (**103**) or 3,3-dimethylbutyryl groups conferred acid stability and a longer antagonism of the CCK-8-induced hypomotility in mice by oral administration. Moreover, compound **103** by intraperitoneal or oral administration showed protective effect on experimental acute pancreatitis induced by caerulein in rats.²⁴⁴ Respecting the N2-substituent, the SAR study indicated the importance of its lipophilic character and an appropriate spatial orientation. Thus, the benzyl group at this position in the prototype **95** was replaced by cyclohexyl, phenyl, or naphthyl groups without affecting binding affinity and selectivity, while its replacement by methyl group led to almost the complete loss of the binding affinity.²⁴⁵ It is interesting to note the existence of atropoisomerism in the 2-naphthyl derivatives, and that, among the two epimers at this position, the (2*S*)-compound **111** was 26-fold more active than its (2*R*)-epimer (IC₅₀ = 15.4 nM). This difference between both epimers shows the importance of the aryl group orientation. In general, the insertion of substituents with different electronic or steric properties into

Table XIV. 1,3-Dioxoperhydropyrido[1,2-*c*]pyrimidine-Based CCK₁ Receptor Antagonists

Compound	IC ₅₀ (nM)		Selectiv. CCK ₂ /CCK ₁	Ref.					
	CCK ₁ ^a	CCK ₂ ^b							
CCK-8	1.04	5.60	5.4	244					
 94	2,350	>10,000	>4	237					
 95	Configuration		K_i (nM)						
	N ^o	4a			5				
	IQM-95,333 (95)	S			R				
	96	R			S				
	97	R			R				
98	S	S	2,890	2,910	1				
 99	N ^o	X	R ¹	R ²	R ³	IC ₅₀ (nM)			
	95	CO	In ^c	Boc	H	1.59	>10,000	6,300	
	99	CO	Ph	Boc	H	65.7	>10,000	152	
	100	CO	In ^c	Boc	Me	42.4	>10,000	236	
	101	CO	In ^c	H	H	>1,000	>10,000	---	244
	102 ^d	CO	In ^c	2-Adc ^e	H	340	3,430	10	
	103	CO	In ^c	tBu-HNCO	H	0.91	>10,000	>11,000	
	104	CH ₂	In ^c	Boc	H	>1,000	>10,000	---	
	105	[(R)CHCN]	In ^c	Boc	H	7.69	>10,000	>1,300	
 106	N ^o	X	Y	R ¹	IC ₅₀ (nM)				
	106	CO	CO	Me	9,162	>10,000	>1	245	
	107	CO	CO	Cyclohexyl	0.60	>10,000	>16,700	245	
	108	CO	CO	Ph	1.18	>10,000	>8,500	245	
	109	CO	CO	(S)-CH(CH ₃)Ph	6.89	>10,000	>1,450	245	
	110	CO	CO	2-(Me)Ph	0.97	>10,000	>10,300	245	
	111	CO	CO	1-Naphtyl	0.59	>10,000	>17,000	245	
	112	CS	CO	CH ₂ Ph	0.09	>10,000	>10 ⁵	246	
	113	CS	CS	CH ₂ Ph	1.34	>10,000	7,500	246	
	114	CO	CS	CH ₂ Ph	2.83	>10,000	3,500	246	
	115	CH ₂	CO	CH ₂ Ph	>1,000	>10,000	---	246	
	116	CO	CH ₂	CH ₂ Ph	>1,000	>10,000	---	246	
 117					>1,000	>10,000	---	246	


^aIC₅₀ or K_i values for the inhibition of the binding of [propionyl-³H]CCK-8 to rat pancreas.^bIC₅₀ or K_i values for the inhibition of the binding of [propionyl-³H]CCK-8 to rat cerebral cortex membranes.^cIn = indol-3-yl.^d(4aR,5S)-stereochemistry.^e2-Adc = 2-adamantylloxycarbonyl.

the phenyl group of compound **108** led to (2–50)-fold reduction in the binding affinity at CCK₁ receptors, independently on the nature of the substituent.²⁴⁵

Modifications at the central 1,3-dioxoperhydropyrido[1,2-*c*]pyrimidine skeleton have indicated that, whereas the replacement of the 1- or/and 3-oxo groups of IQM-95,333 by the thioxo analogues (compounds **112–114**) is allowed, the reduction of these groups (**115** and **116**) or the contraction of the fused piperidine ring to the pyrrolidine analogue (**117**) led to the complete loss of the binding affinity.²⁴⁶ These results gave further support for the crucial influence of the topography defined by the 1,3-dioxoperhydropyrido[1,2-*c*]pyrimidine scaffold, with a (4*aS*,5*R*)-stereochemistry upon the binding to CCK₁ receptors. The 1-thioxo derivative **112**, with subnanomolar affinity (0.09 nM), even higher than that of the endogenous ligand CCK-8, and > 10⁵-fold selectivity over the CCK₂ receptors, is the most selective and one of the most potent CCK₁ antagonists reported up to now.

The high CCK₁ selectivity of the 1,3-dioxoperhydropyrido[1,2-*c*]pyrimidine-based CCK antagonists has recently been reversed towards the CCK₂ receptors by combination of configuration inversion at this bicyclic heterocyclic system, replacement of the Boc group by the 2-adamantyloxycarbonyl, and insertion of a methyl group into the (4*S*)-position, such as in compounds **118** and **119** of Table XV.²⁴⁷

Table XV. 1,3-Dioxoperhydropyrido[1,2-*c*]pyrimidine-Based CCK₂ Receptor Antagonists

	N°	R ¹	-IC ₅₀ (nM)		Selectivity CCK ₁ /CCK ₂	Ref.
			CCK ₁ ^a	CCK ₂ ^b		
	118	CH ₂ Ph	>10,000	181	>83	247
	119	4-[N(CH ₃) ₂]Ph	>10,000	121	>55	247

^aIC₅₀ for the inhibition of the binding of [propionyl-³H]CCK-8 to rat pancreas.






^bIC₅₀ for the inhibition of the binding of [propionyl-³H]CCK-8 to rat cerebral cortex membranes.

J. Pyrazolidinone and Related Heterocyclic-Derived CCK Antagonists

Efforts to optimize lead compounds through random screening used at the Lilly company resulted in the discovery of diphenylpyrazolidinone derivatives with nanomolar affinity at CCK₁ and CCK₂ receptors as compounds LY294920 (Table XVI, **121**, X = S) and LY288513 (**120**, X = O), respectively.²⁴⁸ In the rat stomach, the CCK₂ antagonist LY288513 (**120**) did not inhibit the gastrin-induced activation of histidine decarboxylase.²⁴⁹ However, this compound produced anxiolytic-like effects in mice and rats.^{250–252} In the elevated plus-maze test, this antagonist showed an anxiolytic effect in mice of comparable potency to diazepam, used as a reference standard. However, unlike diazepam, LY288513 did not affect muscle tone, neuromuscular coordination, or sensorimotor reactivity. High doses of this compound were required to reduce spontaneous activity levels, decrease body temperature, or potentiate the CNS-depressant effects of hexobarbital. LY288513 had no analgesic activity in mouse writhing or tail-flick tests. Electrophysiological studies in anaesthetized rats showed that acute administration of LY288513 decreased the number of spontaneously active dopamine neurons in the substantia nigra and ventral tegmental area, but did not produce catalepsy. These results indicated that LY288513 could possess both anxiolytic and antipsychotic potential.²⁵⁰ Moreover, LY288513 blocked the effects of nicotine²⁵³ and diazepam²⁵⁴ withdrawal in rats. However, the development of LY288513 was discontinued because of adverse effects in preclinical toxicological studies.⁷⁸

Researchers at Searle employed 1,3,5-trisubstituted pyrrolidinones as scaffolds for appending groups which could mimic amino acid side chains of CCK-4.²⁵⁵ This approach led to CCK₁ antagonists represented by SC-50,998 (**122**) and **123** (Table XVI). The carboxylic acid of these compounds is essential for binding affinity. SC-50,998 competitively inhibited the CCK-8-stimulated

Table XVI. Pyrazolidinone and Related Heterocyclic-Based CCK Antagonists

Compound	IC ₅₀ (nM)		Selectivity CCK ₁ /CCK ₂	Ref.
	CCK ₁	CCK ₂		
	N ^o	Conf. 4 5	R	
	LY288513 (120)	S R	4-Br	20,500 ^a 19 ^b 1,080 248
	LY294920 (121)	R S	3-CF ₃ 4-Cl	17 ^a 1,900 ^b 0.009 248
	N ^o		R	
	SC-50,998 (122)			16 ^c >10,000 ^d <0.002 255
	123			18.5 ^c 4,800 ^d 0.0038
	N ^o	R ¹	R ²	
	124	H	H	823 ^e 16 ^f 51
	125	3-F	H	1,000 ^e 31 ^f 32 256
	126	H	2-CH ₃	580 ^e 7.8 ^f 74
	127	H	2-CH ₂ CH ₃	810 ^e 25 ^f 32


^aInhibition of the [³H]L-364,718 binding to mouse brain.^bInhibition of the [¹²⁵I]CCK-8 binding to mouse brain.^cEvaluated in rat pancreatic membranes.^dEvaluated in guinea pig brain cortex.^eInhibition of the [¹²⁵I]CCK-8 binding to guinea pig pancreas.^fInhibition of the [¹²⁵I]CCK-8 binding to guinea pig cortex.

contraction of guinea pig ileal smooth muscle. By intraperitoneal and oral administration, this compound reversed the CCK-8-induced delayed gastric emptying in rats.²⁵⁵

The hexahydroazepin-2-one ring, with a similar pattern of substituents to that used in the pyrazolidinone derivatives, was used at Pfizer as an alternative to the benzodiazepine nucleus.²⁵⁶ This approach led to some compounds with nanomolar affinity at CCK₂ receptors, such as **126**. However, no functional or pharmacological activity of these compounds have been reported.

K. Indol-2-one-Based CCK Antagonists

The 1,3,3-trisubstituted indol-2-one derivatives AG-041R (Table XVII, **128**) and T-0632 (**129**) were selected as selective CCK₂ and CCK₁ receptor antagonists, respectively, from random screening programs. AG-041R (**128**) inhibited pentagastrin-stimulated acid secretion by i.v. administration (ID₅₀ = 5 nM/kg) and had no inhibitory effect on carbachol or histamine-stimulated secretion.²⁵⁷ This antagonist exhibited greater potency than L-365,260 in the water-immersion stress (600-fold) and indomethacin-induced ulcer (6-fold) models.²⁵⁸ AG-041R dose-dependently inhibited the gastrin-induced histidine decarboxylase activation in intact fasted rats with a ID₅₀ value of 0.01 μmol/kg i.v., 5-fold lower than that of the benzodiazepine derivative YF476.²³¹ This activity has also been studied in enterochromaffin-like (ECL) carcinoma tumors in *Mastomys natalensis* rats both *in vitro* and *in vivo*, where AG-041R also inhibited the histidine decarboxylase gene expression in ECL carcinoid tumor cells and the gastrin-induced DNA synthesis and *c-fos* gene expression.²⁵⁹

Table XVII. Indol-2-One-Based CCK Antagonists

Compound	Affinity		Selectivity CCK ₁ /CCK ₂	Reference
	CCK ₁	CCK ₂		
AG-041R (128)	555 ^a	1.11 ^b	500	83
T-0632 (129)	0.24 ^c	5,600 ^d	4 × 10 ⁻⁵	263

^aIC₅₀ (nM) value of inhibition of the binding of [¹²⁵I]CCK-8 to guinea pig pancreas.

^bIC₅₀ (nM) value of inhibition of the binding of [¹²⁵I]gastrin to guinea pig gastric glands.

^cK_i (nM) value of inhibition of the binding of [¹²⁵I]CCK-8 to rat pancreas.

^dK_i (nM) value of inhibition of the binding of [¹²⁵I]CCK-8 to guinea pig cortex.


Additionally, AG-041R induced systemic cartilage hyperplasia by stimulation of chondrocyte proliferation and metabolism, an intrinsic property of this compound not related with its CCK₂ receptor antagonism.^{260,261} AG-041R also stimulated the repair of osteochondral defects in a rabbit model.²⁶² These findings suggest that this compound could be a therapeutic agent for cartilage disorders.

As shown in Table XVII, T-0632 (129) is a potent and highly selective CCK₁ receptor antagonist, which inhibited the CCK-8-stimulated amylase release from rat pancreatic acini in a concentration-dependent manner with an IC₅₀ value of 5.0 nM. This compound also competitively inhibited the CCK-8-induced contraction of the rabbit gall bladder smooth muscle with a pA₂ value of 8.5.²⁶³ In rats, by intravenous and intraduodenal routes, T-0632 dose-dependently inhibited the CCK-8-stimulated pancreatic secretion with ED₅₀ values of 0.025 and 0.040 mg/kg, respectively. In mice, orally administered, T-0632 prevented the caerulein-induced pancreatitis, the CCK-8-induced inhibition of gastric emptying, and the CCK-8-induced gall bladder emptying in a dose-dependent manner with ED₅₀ values of 0.028, 0.04, and 0.12 mg/kg, respectively. In dogs T-0632 inhibited CCK-8-stimulated pancreatic amylase secretion at a dose of 0.01 mg/kg. In this study, the effects of this CCK₁ antagonist were more selective for the pancreas than for the gall bladder.²⁶⁴ In different models of experimental acute pancreatitis, T-0632 has shown preventive effects by oral or intraduodenal administration,^{265,266} and has been studied in Phase I clinical trials for the treatment of pancreatic disorders, although results of these studies have not been reported.

L. Other CCK Antagonists

1. Lintitript (SR-27,897)

The CCK₁ selective antagonist SR-27,897 (also designed with the generic name of Lintitript, compound 130 of Table XVIII) was obtained by optimisation of a lead compound discovered through random screening of a large chemical library at Sanofi.²⁶⁷ This compound competitively inhibited the CCK-8-stimulated amylase release in isolated rat pancreatic acini (pA₂ = 7.50) and the CCK-8-induced guinea pig bladder contractions (pA₂ = 9.57). This antagonistic activity was confirmed in *in vivo* gastrointestinal models. Thus, at 1 mg/kg (i.v.) it completely reversed the CCK-induced amylase secretion in rats, and 3 µg/kg (p.o.) antagonized by 50% the CCK-induced inhibition of gastric emptying in mice, and inhibited the CCK-induced gall bladder emptying in mice with a ED₅₀

Table XVIII. Miscellaneous Structure CCK Antagonists

Compound	Affinity		Selectivity CCK ₂ /CCK ₁	Reference
	CCK ₁	CCK ₂		
SR-27,897 (130) (Lintitript)	0.58 ^a	489 ^b	843	267
TP-680 (131)	1.2 ^c	1,812 ^d	1,510	278
Tetronothiodin (132)	>100,000 ^a	3.6 ^e	<4×10 ⁻⁵	280

^aIC₅₀ (nM) value of inhibition of the binding of [¹²⁵I]CCK-8 to rat pancreas.^bIC₅₀ (nM) value of inhibition of the binding of [¹²⁵I]CCK-8 to guinea pig cortex.^cK_i (nM) value of inhibition of the binding of [¹²⁵I]CCK-8 to rat pancreas.^dK_i (nM) value of inhibition of the binding of [¹²⁵I]CCK-8 to rat cerebral cortex.^eIC₅₀ (nM) value of inhibition of the binding of [¹²⁵I]CCK-8 to rat cerebral cortex.

value of 27 µg/kg (p.o.). SR-27,897 showed a long-lasting action in all the experiments, with no differences between oral and intravenous routes of administration. In comparison with devazepide, SR-27,897 increased the gall bladder volume of fasting mice, but its effect was 10-fold lower than that of devazepide.²⁶⁷ SR-27,897 dose-dependently antagonized the CCK-8-induced hypophagia and hypolocomotion in rats, two behavioural effects associated with the stimulation of CCK₁ receptors, with ED₅₀ values of 0.003 and 0.002 mg/kg (i.p.), respectively, while devazepide in the same tests showed ED₅₀ values of 0.02 and 0.1 mg/kg (i.p.), respectively. In these tests the i.p./p.o. ratio for SR-27,897 was near unity, suggesting a high oral bioavailability.²⁶⁸ SR-27,897 inhibited the preprandial pancreatic secretion and greatly reduced the postprandial pancreatic juice and enzyme outflows in milk-fed calves, demonstrating the implication of CCK and CCK₁ receptors in mediating the postfeeding pancreatic response.²⁶⁹ Similarly, the inhibitory effect of SR-27,897 on the CCK-8-induced contraction of human lower oesophageal sphincter strips demonstrated that the contractile effect of CCK is mainly due to the activation of CCK₁ receptors.²⁷⁰ SR-27,897 also inhibited the pressor effects of CCK on blood pressure and reversed the bradycardic responses to tachycardia in conscious rats, providing evidence for the involvement of CCK₁ receptors in cardiovascular regulation.²⁷¹ On the other hand, neither SR-27,897 nor devazepide exhibited anxiolytic-like effects in the elevated zero-maze test in rats, while the CCK₂ antagonists L-365,260 and PD-135,158 both had significant anxiolytic activity in this assay.²⁷² In Phase II clinical studies, SR-27,897 accelerated gastric emptying of solids, while gastric emptying of liquids was not significantly altered by oral administration in healthy male volunteers. Moreover, this CCK₁ antagonist markedly increased postprandial plasma CCK release, while distinctly reduced postprandial pancreatic polypeptide.²⁷³ Several receptor mutagenesis studies have shown that SR-27,897 occupies different binding sites from its analogue CCK₁ agonist SR-146,131 and CCK-8 at the CCK₁ receptor.²⁷⁴⁻²⁷⁷

2. TP-680

The selective CCK₁ receptor antagonist TP-680 (Table XVIII, 131) was also the result of a random screening program. This antagonist was approximately 17-fold less potent than devazepide, but 106-

fold more potent than loxiglumide in inhibiting the CCK-8-stimulated amylase release from rat pancreatic acini.²⁷⁸ TP-680 by intravenous administration in mice has shown long-lasting antagonistic properties on CCK-8-stimulated pancreatic secretion, gastric emptying, and gall bladder contraction. The selectivity for these activities was gastric emptying > pancreatic secretion > gall bladder contraction.²⁷⁹

3. Tetronothiodin

The highly selective CCK₂ antagonist tetronothiodin (**132**) was isolated from the fermentation broth of *Streptomyces* sp. NR0489.^{280,281} This compound inhibited the CCK-8 binding to rat cerebral cortex membranes with nanomolar binding affinity, whereas it did not show affinity at rat pancreatic membranes. It also inhibited the CCK-8-induced Ca²⁺ mobilization in rat anterior pituitary cells GH3, but it did not inhibit the basal cytosolic Ca²⁺ concentration.²⁸⁰

4. CONCLUSIONS AND FUTURE PERSPECTIVES

Over the past 15 years, the search of CCK receptor ligands has evolved from the initial CCK structure derived peptides towards peptidomimetic or non-peptide agonists and antagonists with improved pharmacokinetic profile. This research has provided a broad assortment of potent and highly selective antagonists for both CCK receptor subtypes, CCK₁ and CCK₂, of diverse chemical structure. These antagonists, as pharmacological tools, have highly contributed to the characterization and localization of CCK receptor subtypes, as well as to the study of physiological and pathological actions of CCK. However, despite the progress in this field, the complex biological effects of CCK mediated by CCK₁ and CCK₂ receptors are not yet completely established. Particularly, additional research is necessary for gaining insight into the complex system of interaction of CCK with other neurotransmitters both in the CNS and in the periphery. Pharmacological research is also necessary to confirm that the different binding affinities determined for some antagonists by using different radioligands, and in some cases the discrepancies observed between binding potency and antagonistic potency, are due to the existence of different binding states and not to receptor subtype heterogeneity. This research will help to characterize the different binding states for each receptor subtype as well as the biological actions mediated by their activation. Furthermore, at the molecular level, as in the case of other G-protein coupled receptors, the growing studies focused on the characterization of agonists and antagonists binding domains at both CCK₁ and CCK₂ receptors, would contribute to a better knowledge of the complex dynamic equilibrium in the receptor activation-inactivation process and their related biological responses. Additionally, these molecular studies will foster the *de novo* receptor structure-based design of new selective and more effective antagonists.

Concerning the therapeutic potential of CCK antagonists, although several CCK₁ and CCK₂ receptor antagonists, summarized in Table XIX, have reached different phases of clinical trials, the complex interconnected physiological actions of CCK have hampered their clinical development. On one hand, CCK₁ antagonists entered development mainly for the treatment of pancreatic disorders and as prokinetics for the treatment gastroesophageal reflux disease, bowel disorders (e.g., irritable bowel syndrome), and gastroparesis. Their potential in the treatment of pancreatitis and pancreatic cancer is showing particular promise. However, their therapeutic development has been hampered by the side effect of stone growth in the gall bladder, which was firstly reported from the clinical trial of devazepide (**19**). Conversely, this side effect has not been reported for loxiglumide (**7**), the most advanced CCK₁ antagonist in clinical studies for the treatment of acute pancreatitis, or for lintitript (**130**). If this side effect were confirmed as a general property related to the CCK₁ receptor antagonism, those antagonists as T-0632 (**129**) which are more selective for the pancreas than for the gall bladder would have some advantage.

Table XIX. Summary of CCK Antagonists That Have Reached Clinical Trials

<i>Compound</i>	<i>Receptor subtype selectivity</i>	<i>Highest clinical phase^a</i>	<i>Indication</i>
Proglumide (3)	Non-selective	Marketed	Antiulcer
Loxiglumide (7)	CCK ₁	Pre-registered	pancreatitis
Dexloxiglumide (8)	CCK ₁	Phase III	Irritable bowel syndrome
KSG-504 (11)	CCK ₁	Phase I	Pancreatic disorders
2-NAP (12)	CCK ₁	Phase I	Pancreatic disorders
Spiroglumide (13)	CCK ₂	Phase II	Gastric secretion disorders
Itriglumide (17)	CCK ₂	Phase I	Anxiolytic, antiulcer
Devazepide (18)	CCK ₁	Phase II (discontinued)	Gastric motility disorders
		Phase II	Pain
Pranazepide (20)	CCK ₁	Phase II	Pancreatic disorders
Tarazepide (21)	CCK ₁	Phase II	Gastric emptying disorders
L-365,260 (22)	CCK ₂	Phase II (discontinued)	Anxiety, drug dependence
		Phase II	Pain
YM022 (30)	CCK ₂	Phase I	Gastric secretion disorders
YF476 (31)	CCK ₂	Phase I	Gastro-oesophageal reflux
GV150013X (41)	CCK ₂	Phase II	Anxiety disorders
		Phase II	Sleep disorders
S-0509 (49)	CCK ₂	Phase I	Gastric secretion disorders
CI-988 (58)	CCK ₂	Phase II	Anxiety and panic disorders
T-0632 (129)	CCK ₁	Phase I	Pancreatic disorders
Lintitript (130)	CCK ₁	Phase II	Pancreatic cancer therapy, eating disorders

^aInformation from the Ensemble Data Base of Prous Science.

On the other hand, CCK₂ antagonists have been entered development mainly for the treatment of gastric acid secretion and anxiety disorders. Although CCK₂ receptor antagonists represent an alternative therapeutic approach for the treatment of peptic ulcer and gastric acid secretion disorders, the clinical results of L-365,260 (22) and spiroglumide (13) are not encouraging as relatively high doses were required to obtain results equivalents to those of the H₂-receptor antagonist famotidine or the proton pump inhibitor omeprazole. The effectiveness of the current treatments of duodenal ulcer patients through the use of these H₂-receptor antagonists or proton pump inhibitors supplemented by *H. pylori* eradication has diminished the therapeutic opportunity for CCK₂ antagonists in the therapy of gastric acid-related diseases. However, the therapeutic potential of these antagonists will be fully evaluated only when the results of the in-progress clinical trials of the second generation CCK₂ antagonists, with improved oral bioavailability, such as itriglumide (17) and YF476 (30), are available. Concerning this issue, it is important to note that, in *in vivo* animal models, itriglumide was effective in prevention of gastric damage and YF476 reversed the hypergastrinemia and cell proliferation caused by omeprazole in the gastric mucosa.

The discovery of the panicogenic effect of CCK-4 in man raised the hypothesis of the involvement of CCK₂ receptors in the pathogenesis of panic disorders, consequently CCK₂ antagonists were considered potential anxiolytic agents. In fact, most of the potent CCK₂ antagonists have shown anxiolytic-like effects in diverse animal models, without the side effects of the classic benzodiazepine anxiolytics, such as sedation, development of tolerance, and withdrawal anxiogenesis after termination of the treatment. However, these anxiolytic effects have not been confirmed in clinical trials neither in patients with generalized anxiety and panic disorders neither against CCK-4-induced panic symptoms in healthy volunteers. These discouraging clinical results have been attributed to the poor bioavailability and blood-brain permeability of the first generation CCK₂ antagonists studied [L-365,260 (22) and CI-988 (58)], and it has been suggested that the second generation antagonists with improved oral bioavailability might give more encouraging results.

Nevertheless, there are doubts about the actual role of CCK₂ receptors in anxiogenesis.¹¹ Regarding this issue, it is noteworthy that recent studies have shown that CCK₂ receptor-deficient mice did not show behavioral modifications compared to wild-type mice in the elevated plus maze test of anxiety and in the motility conditioned suppression test, indicating that compensatory mechanisms very likely occur following CCK₂ receptor inactivation.²⁸² Therefore, considerable pharmacological work is still needed to really assess the therapeutic potential of both CCK₁ and CCK₂ receptor antagonists.

ACKNOWLEDGMENTS

The author thanks Dr. M. Teresa García-López for reading the manuscript and making valuable suggestions.

REFERENCES

- Ivy AC, Oldberg E. A Hormone mechanism for gall bladder contraction and evacuation. *Am J Physiol* 1928;86:599–613.
- Jorpes E, Mutt V. Cholecystokinin and pancreozymin, one single hormone? *Acta Physiol Scand* 1966;66:196–202.
- Mutt V, Jorpes JE. Structure of porcine cholecystokinin-pancreozymin. 1. Cleavage with thrombin and with trypsin. *Eur J Biochem* 1968;6:156–162.
- Vanderhaeghen JJ, Signeau JC, Gepts W. New peptide in vertebrate CNS reacting with antigastrin antibodies. *Nature* 1975;267:604–605.
- Beinfeld MC, Meyer DK, Eskay RL, Jensen JT, Brownstein MJ. The distribution of cholecystokinin-immunoreactivity in the central nervous system of the rat, as determined by radioimmunoassay. *Brain Res* 1981;212:51–57.
- Beinfeld MC, Palkovits M. Distribution of cholecystokinin (CCK) in the rat lower brain stem nuclei. *Brain Res* 1982;238:260–265.
- Hökfelt T, Skirboll L, Everitt B, Meister B, Brownstein M, Jacobs T, Faden A, Kuga S, Goldstein M, Markstein R, Döckray G, Rehfeld J. Distribution of cholecystokinin-like immunoreactivity in the nervous system: Co-existence with classical neurotransmitters and other neuropeptides. *Ann NY Acad Sci* 1985;448:255–274.
- Crawley JN, Corwin RL. Biological actions of cholecystokinin. *Peptides* 1994;15:731–755.
- Daugé V, Léna I. CCK in anxiety and cognitive processes. *Neurosci Biobehav R* 1998;22:815–825.
- Griebel G. Is there a future for neuropeptide receptor ligands in the treatment of anxiety disorders? *Pharmacol Ther* 1999;82:1–61.
- Rehfeld JF. Cholecystokinin and panic disorder—three unsettled questions. *Regul Peptides* 2000;93:79–83.
- Ritter RC, Covasa M, Matson CA. Cholecystokinin: Proofs and prospects for involvement in control of food intake and body weight. *Neuropeptides* 1999;33:387–399.
- Moran TH. Cholecystokinin and satiety: Current perspectives. *Nutrition* 2000;16:858–865.
- Degen L, Matzinger D, Drewe J, Beglinger C. The effect of cholecystokinin in controlling appetite and food intake in humans. *Peptides* 2001;22:1265–1269.
- Wiesenfeld-Hallin Z, Xu X-J. The role of cholecystokinin in nociception, neuropathic pain and opiate tolerance. *Regul Peptides* 1996;65:23–28.
- Szelényi Z. Cholecystokinin and thermoregulation—A minireview. *Peptides* 2001;22:1245–1250.
- Itoh S, Lai H. Influences of cholecystokinin and analogues on memory processes. *Drug Dev Res* 1990;21:257–276.
- Noble F, Roques BP. CCK-B Receptor: Chemistry, molecular biology, biochemistry and pharmacology. *Prog Neurobiol* 1999;58:349–379.
- Ghijzen WEJM, Leenders AGM, Wiegant VM. Regulation of cholecystokinin release from central nerve terminals. *Peptides* 2001;22:1213–1221.
- Crawley JN. Cholecystokinin-dopamine interactions. *Trends Pharmacol Sci* 1991;12:232–236.
- Tanganelli S, Fuxe K, Antonelli T, O'Connor WT, Ferraro L. Cholecystokinin/dopamine/GABA interactions in the nucleus accumbens: Biochemical and functional correlates. *Peptides* 2001;22:1229–1234.
- Shlik J, Vasar E, Bradwejn J. Cholecystokinin and psychiatric disorders. *CNS Drugs* 1997;8:134–152.

23. Noble F, Wank SA, Crawley JN, Bradwejn J, Seroogy KB, Hamon M, Roques BP. International Union of Pharmacology. XXI. Structure, distribution, and functions of cholecystokinin receptors. *Pharmacol Rev* 1999;51:745-781.
24. Crespi F, Corsi M, Reggiani A, Ratti E, Gaviraghi G. Involvement of cholecystokinin within craving for cocaine: Role of cholecystokinin receptor ligands. *Exp Opin Invest Drugs* 2000;9:2249-2258.
25. Deschenes RJ, Lorenz LJ, Haun RS, Roos BA, Collier KJ, Dixon JE. Cloning and sequence analysis of cDNA encoding rat preprocholecystokinin. *Proc Natl Acad Sci USA* 1984;81:726-730.
26. Martinez J. Gastrointestinal regulatory peptide receptors. In Hansch C, Sammes PG, Taylor JB, editors. *Comprehensive Medicinal Chemistry*. Vol 3, Emmett JC, editor. Oxford: Pergamon Press; 1990. p 926-959.
27. Wank SA. Cholecystokinin receptors. *Am J Physiol* 1995;269:G628-G646.
28. Vanhoutte PM, Humprey PPA, Spedding M. XI. International Union of Pharmacology. Recommendations for nomenclature of new receptor subtypes. *Pharmacol Rev* 1996;48:1-2.
29. Wank SA. G Protein-coupled receptors in gastrointestinal physiology I. CCK receptors: An exemplary family. *Am J Physiol* 1998;274:G607-G613.
30. Baber NS, Dourish CT, Hill DR. The role of CCK, caerulein and CCK antagonists in nociception. *Pain* 1989;39:307-328.
31. Feliciano LF, Mazzini BK, Cacheiro RGR, Cruz TN, Flórido JC, Nasello AG. Stimulation of either cholecystokinin receptor subtype reduces while antagonists potentiate or sensitize a morphine-induced excitatory response. *Peptides* 2001;22:1299-1304.
32. Logsdon CD. Glucocorticoids increase cholecystokinin receptors and amylase secretion in pancreatic acinar AR42J cells. *J Biol Chem* 1986;261:2096-2101.
33. Klueppelberg UG, Molero X, Barrett RW, Miller LJ. Biochemical characterization of the cholecystokinin receptor on CHP212 human neuroblastoma cells. *Mol Pharmacol* 1990;38:159-163.
34. Wank SA, Harkins RT, Jensen RT, Shapira H, de Weerth A, Slattery T. Purification, molecular cloning, and functional expression of the cholecystokinin receptor from rat pancreas. *Proc Natl Acad Sci USA* 1992;89:3125-3129.
35. de Weerth A, Pisegna JR, Wank S. Guinea pig gall bladder and pancreas possess identical CCK-A receptor subtypes: Receptor cloning and expression. *Am J Physiol* 1993;265:G1116-G1121.
36. de Weerth A, Pisegna JR, Huppi IK, Wank S. Molecular cloning, functional expression and chromosomal localization of the human cholecystokinin type A receptor. *Biochem Biophys Res Commun* 1993;194:811-818.
37. Ulrich CD, Ferber I, Holicky E, Hadac E, Buell G, Miller LJ. Molecular cloning and functional expression of the human gall bladder cholecystokinin A receptor. *Biochem Biophys Res Commun* 1993;193:204-211.
38. Reuben M, Rising L, Prinz C, Hersey S, Sachs G. Cloning and expression of the rabbit gastric CCK-A receptor. *Biochim Biophys Acta* 1994;1219:321-327.
39. Tatsuura G, Shinohara S, Shintaku H, Eigyo M, Matsushita A. Protective effect of CCK-8 and ceruletide on glutamate-induced neuronal cell death in rat neuron cultures: Possible involvement of CCK-B receptors. *Neurosci Lett* 1991;132:159-162.
40. Yasui M, Kawasaki K. CCK-B receptor activation protect CA1 neurons from ischemia-induced disfunction in stroke-prone spontaneously hypertensive rats hippocampal slices. *Neurosci Lett* 1995;191:99-102.
41. Eigyo M, Katsuura G, Shintaku H, Shinohara S, Shiomi T, Matsushita A. Systemic administration of a cholecystokinin analogue, ceruletide protects against ischemia-induced neurodegeneration in gerbils. *Eur J Pharmacol* 1992;214:149-158.
42. Sacerdote P, Wiedermann CJ, Wahl LM, Pert CB, Ruff MR. Visualization of cholecystokinin receptors on a subset of human monocytes and in rat spleen. *Peptides* 1991;12:167-176.
43. Lignon MF, Bernad N, Martinez J. Pharmacological characterization of type B cholecystokinin binding sites on the human JURKAT T lymphocyte cell line. *Mol Pharmacol* 1991;39:615-620.
44. Rozengurt E, Walsh JH. Gastrin CCK, signaling, and cancer. *Annu Rev Physiol* 2001;63:49-76.
45. Wank SA, Pisegna JR, de Weerth A. Brain and gastrointestinal cholecystokinin receptor family: Structure and functional expression. *Proc Natl Acad Sci USA* 1992;89:8691-8695.
46. Menozzi D, Gardner JD, Maton PN. Properties of receptors for gastrin and CCK on gastric smooth muscle cells. *Am J Physiol* 1989;257:G73-G79.
47. Pisegna JR, de Weerth A, Wank SA. Molecular cloning of the human brain and gastric cholecystokinin receptor: Structure, functional expression and chromosomal localization. *Biochem Biophys Res Commun* 1992;189:296-303.
48. Kopin AS, Lee Y-M, McBride EW, Miller LJ, Lu M, Lin HY, Kolakowski LF, Beinborn M. Expression cloning and characterization of the canine parietal cell gastrin receptor. *Proc Natl Acad Sci USA* 1992;89:3605-3609.

49. de Weerth A, von Schrenck T, Gronewold M, Freudenberg F, Mirau S, Schulz M, Greten H. Characterization of CCK receptors in stomach smooth muscle: Evidence for two subtypes. *Biochim Biophys Acta* 1997;1327:213–221.
50. Baldwin GS. Gastrin receptor structure. In: Walsh JH, editor. *Gastrin*. New York: Raven; 1993. p195–207.
51. Bold RJ, Ishizuka J, Townsend CM, Thompson JC. Gastrin stimulates growth of human colon cancer cells via a receptor other than CCK-A or CCK-B. *Biochim Biophys Res Commun* 1994;202:1222–1226.
52. Ishizuka J, Townsend CM, Bold RJ, Martinez J, Rodriguez M, Thompson JC. Effects of gastrin on 3',5'-cyclic adenosine monophosphate, intracellular calcium, and phosphatidylinositol hydrolysis in human colon cancer cells. *Cancer Res* 1994;54:2129–2135.
53. Baldwin GS. Antiproliferative gastrin/cholecystokinin receptor antagonists target the 78-kDa gastrin-binding protein. *Proc Natl Acad Sci USA* 1994;91:7593–7597.
54. Seva C, Dickinson CJ, Yamada T. Growth-promoting effects of glycine-extended progastrin. *Science* 1994;265:410–412.
55. Alexander SPH, Mathie A, Peters AJ, editors. *Cholecystokinin and gastrin receptors*. Trends Pharmacol Sci, 2001 Nomenclature Supplement 2001;22:36–37.
56. Kopin AS, McBride EW, Quinn SM, Kolakowski LF, Beinborn M. The role of the cholecystokinin-B/gastrin receptor transmembrane domains in determining affinity for subtype-selective ligands. *J Biol Chem* 1995;270:5019–5023.
57. Silvente-Poirot S, Wank SA. A segment of five amino acids in the second extracellular loop of the cholecystokinin-B receptor is essential for selectivity of the peptide agonist gastrin. *J Biol Chem* 1996;271:14698–14706.
58. Silvente-Poirot S, Escrieut C, Wank SA. Role of the extracellular domains of the cholecystokinin receptor in agonist binding. *Mol Pharmacol* 1998;54:364–371.
59. Beinborn M, Lee Y-M, McBride EW, Quinn SM, Kopin AS. A single amino acid of cholecystokinin-B/gastrin receptor determines specificity for non-peptide antagonists. *Nature* 1993;362:348–350.
60. Ding X-Q, Miller LJ. Characterization of the type A cholecystokinin receptor hormone-binding domain: Use of contrasting and complementary methodologies. *Peptides* 2001;22:1223–1228.
61. Bläker M, Ren Y, Gordon MC, Hsu JE, Beinborn M, Kopin AS. Mutations within the cholecystokinin-B/gastrin receptor ligand 'Pocket' interconvert the functions of nonpeptide agonists and antagonists. *Mol Pharmacol* 1998;54:857–863.
62. Kopin AS, McBride EW, Schaffer K, Beinborn M. CCK receptor polymorphisms: An illustration of emerging themes in pharmacogenomics. *Trends Pharmacol Sci* 2000;21:346–352.
63. Tachikawa H, Harada S, Kawanishi Y, Okubo T, Shiraishi H. Novel polymorphism in the promoter and coding regions of the human cholecystokinin B receptor gene: An association analysis with schizophrenia. *Am J Med Genet* 1999;88:700–7004.
64. Fujii C, Harada S, Ohkoshi N, Hayashi A, Yoshizawa K, Ishizuka C, Nakamura T. Association between polymorphism of the cholecystokinin gene and idiopathic Parkinson's disease. *Clin Genet* 1999;56:394–399.
65. Funakoshi A, Miyasaka K, Matsumoto H, Yamamori S, Takiguchi S, Kataoka K, Takata Y, Matsusue K, Kono A, Shimokata H. Gene structure of human cholecystokinin (CCK) type-A receptor: Body fat content is related to CCK type-A receptor promoter polymorphism. *FEBS Lett* 2000;466:264–266.
66. Talkad VD, Fortune KP, Pollo DA, Shah GN, Wank SA, Gardner JD. Direct demonstration of three different states of the pancreatic cholecystokinin receptor. *Proc Natl Acad Sci USA* 1994;91:1868–1872.
67. Li Y, Zhu J, Owyang C. Electrical physiological evidence for high- and low-affinity vagal CCK-A receptors. *Am J Physiol* 1999;277:G469–G477.
68. Harper EA, Roberts SP, Shankley NP, Black JW. Analysis of variations in L-365,260 competition curves in radioligand binding assays. *Br J Pharmacol* 1996;118:1717–1726.
69. Harper EA, Shankley NP, Black JW. Characterization of the binding of a novel radioligand to CCKB/gastrin receptors in membranes from rat cerebral cortex. *Br J Pharmacol* 1999;126:1504–1512.
70. Harper EA, Griffin EP, Shankley NP, Black JW. Analysis of the behaviour of selected CCKB/gastrin receptor antagonists in radioligand binding assays performed in mouse and rat cerebral cortex. *Br J Pharmacol* 1999;126:1496–1503.
71. Derrien M, McCort-Tranchepain I, Ducos B, Roques BP, Durieux C. Heterogeneity of CCK-B receptors involved in animal models of anxiety. *Pharmacol Biochem Behav* 1994;49:133–141.
72. Léna I, Simon H, Roques BP, Dugé V. Opposing effects of two CCK-B agonists on the retrieval phase of a two-trial memory task after systemic injection in the rat. *Neuropharmacol* 1999;38:543–553.
73. Freidinger RM. Cholecystokinin and gastrin antagonists. *Med Res Rev* 1989;9:271–290.
74. Woodruff GN, Hughes J. Cholecystokinin antagonists. *Annu Rev Pharmacol Toxicol* 1991;31:469–501.

75. Nadzan AM, Kerwin JF. Cholecystokinin agonists and antagonists. *Annual Reports in Medicinal Chemistry* 1991;26:191–201.
76. Wettstein JG, Bueno L, Junien JL. CCK antagonists: Pharmacology and therapeutic interest. *Pharmac Ther* 1994;62:267–182.
77. D'Amato M, Rovati LC. Cholecystokinin-A receptor antagonists: Therapies for gastrointestinal disorders. *Exp Opin Invest Drugs* 1997;6:819–836.
78. Makovec F, D'Amato M. CCK_B/gastrin receptor antagonists as potential drugs for peptic ulcer therapy. *Dug Discov Today* 1997;2:283–293.
79. Dunlop J. CCK receptor antagonists. *Gen Pharmac* 1998;4:519–524.
80. de Tullio P, Delarge J, Pirotte B. Recent advances in the chemistry of cholecystokinin receptor ligands (agonists and antagonists). *Curr Med Chem* 1999;6:433–455.
81. Chambers MS, Fletcher SR. CCK-B antagonists in the control of anxiety and gastric acid secretion. In: King FD, Oxford AW, editors. *Progress in Medicinal Chemistry*, Vol 37. Oxford: Elsevier Science; 2000. p 45–81.
82. de Tullio P, Delarge J, Pirotte B. Therapeutic and chemical developments of cholecystokinin receptor ligands. *Exp Opin Invest Drugs* 2000;9:129–146.
83. McDonald IM. CCK₂receptor antagonists. *Exp Opin Ther Patents* 2001;11:445–462.
84. Jensen RT, Jones AW, Gardner JD. COOH-Terminal fragments of cholecystokinin. A new class of cholecystokinin receptor antagonists. *Biochim Biophys Acta* 1983;757:250–258.
85. Martinez J, Magous R, Lignon MF, Laur J, Castro B, Bali JP. Synthesis and biological activity of new peptide segments of gastrin exhibiting gastrin antagonist property. *J Med Chem* 1984;27:1597–1601.
86. Sparkel M, Martinez J, Briet C, Jensen RT, Gardner JD. Cholecystokinin-27-32-amide, a member of a new class of cholecystokinin receptor antagonists. *J Biol Chem* 1983;258:6746–6749.
87. Laur J, Rodriguez M, Aumelas A, Bali JP, Martinez J. Synthesis of analogues of the Des-Phe-NH₂ C-terminal hexapeptide of cholecystokinin showing gastrin antagonist activity. *Int J Peptide Protein Res* 1986;27:386–393.
88. Gonzalez-Muñiz R, Bergeron F, Marseigne I, Durieux C, Roques BP. Boc-Trp-Orn(Z)-Asp-NH₂ and derivatives: A new family of CCK antagonists. *J Med Chem* 1990;33:3199–3204.
89. Lignon MF, Galas MC, Rodriguez M, Laur J, Aumelas A, Martinez J. A synthetic peptide derivative that is a cholecystokinin receptor antagonist. *J Biol Chem* 1987;262:7226–7231.
90. Boomgaarden M, Henklein P, Morgenstern R, Sohr R, Ott T, Martinez J. Synthesis and biological activity of CCK peptides with antagonist activity for CCK-A-receptor. *Eur J Med Chem* 1992;27:955–959.
91. Corringier PJ, Weng JH, Ducos B, Durieux C, Boudeau P, Bohme A, Roques BP. CCK-B Agonist or antagonist activity of structurally hindered and peptidase-resistant Boc-CCK₄ derivatives. *J Med Chem* 1993;36:166–172.
92. Martinez J, Bali JP, Rodriguez M, Castro B, Magous R, Laur J. Synthesis and biological activities of some pseudo-peptide analogues of tetragastrin: The importance of the peptide backbone. *J Med Chem* 1985;28:1874–1879.
93. Herrero S, Suárez-Gea ML, González-Muñiz R, García-López MT, Herranz R, Ballaz S, Barber A, Fortuño A, Del Río J. Pseudopeptide CCK-4 analogues incorporating the ψ [CH(CN)NH] peptide bond surrogate. *Bioorg. Med Chem Lett* 1997;7:855–860.
94. Mendre C, Rodriguez M, Lignon MF, Galas MC, Gueudet C, Worms P, Martinez J. Pharmacological activity of cholecystokinin analogues modified in the Met²⁸-Gly²⁹ region. *Eur J Pharmacol* 1990;186:213–222.
95. Goëtz MA, Lopez M, Monaghan RL, Chang RSL, Lotti VJ, Chen TB. Asperlicin, a novel non-peptidic cholecystokinin antagonist from *Aspergillus alliaceus*. Fermentation, isolation and biological properties. *J Antibiot* 1985;38:1633–1637.
96. Chang RS, Lotti VJ, Monaghan RL, Birnbaum J, Stapley EO, Goetz MA, Albers-Schonberg G, Patchett AA, Liesch JM, Hensens OD, Springer JP. A potent nonpeptide cholecystokinin antagonist selective for peripheral tissues isolated from *Aspergillus alliaceus*. *Science* 1985;230:177–179.
97. Rovati AL, Casula PL, Da Re G. Pharmacological and experimental toxicology of a new non-anticholinergic product with antisecretory and gastroprotective activity (CR 242, xylamide-Milid). *Minerva Med* 1967;58:3656–3670.
98. Rovati AL. Inhibition of gastric secretion by anti-gastrinic and H₂-blocking agents. *Scand J Gastroenterol Suppl* 1976;42:113–118.
99. Hahne WF, Jensen RT, Lemp GF, Gardner JD. Proglumide and benzotript: Members of a different class of cholecystokinin receptor antagonists. *Proc Natl Acad Sci USA* 1981;78:6304–6308.
100. Jensen RT, Jones SW, Gardner JD. Structure-function studies of N-acyl derivatives of tryptophan that function as specific cholecystokinin receptor antagonists. *Biochim Biophys Acta* 1983;761:269–277.

101. Maton PN, Sutliff VE, Jensen RT, Gardner JD. Carbobenzoxy amino acids: Structural requirements for cholecystokinin receptor antagonist activity. *Am J Physiol.* 1985;248:G479–G484.
102. Makovec F, Chisté R, Bani M, Revel L, Stnikar I, Rovati AL. New glutamic and aspartic derivatives with potent CCK-antagonistic activity. *Eur J Med Chem* 1986;21:9–20.
103. Makovec F, Bani M, Cereda R, Chisté R, Pacini MA, Revel L, Rovati LA, Rovati LC. Pharmacological properties of lorglumide as a member of a new class of cholecystokinin antagonists. *Arzneimittelforschung* 1987;37:1265–1268.
104. Setnikar I, Bani M, Cereda R, Chisté R, Makovec F, Pacini MA, Revel L, Rovati LC, Rovati LA. Pharmacological characterization of a new potent and specific nonpolypeptide cholecystokinin antagonist. *Arzneimittelforschung* 1987;37:703–707.
105. Setnikar I, Bani M, Cereda R, Chisté R, Makovec F, Pacini MA, Revel L. Anticholecystokinin activities of loxiglumide. *Arzneimittelforschung* 1987;37:1168–1171.
106. Setnikar I, Chisté R, Makovec F, Rovati LC, Warrington SJ. Pharmacokinetics of loxiglumide after single intravenous or oral doses in man. *Arzneimittelforschung* 1988;38:716–720.
107. Setnikar I, Bani M, Cereda R, Chisté R, Makovec F, Pacini MA, Revel L. Loxiglumide protects against experimental pancreatitis. 1987;37:1172–1174.
108. Ochi K, Harada H, Satake K. Clinical evaluation of cholecystokinin-A-receptor antagonist (loxiglumide) for the treatment of acute pancreatitis. A preliminary clinical trial. Study Group of Loxiglumide in Japan. *Digestion* 1999;60(Suppl 1):81–85.
109. Gielkens HA, de Boer SY, Lam WF, Rovati LC, Lamers CB, Masclee AA. The role of cholecystokinin and the cholinergic system in intravenous amino acid-induced gall bladder emptying. *Eur J Gastroenterol Hepatol* 1997;9:1227–1231.
110. Schwizer W, Borovicka J, Kunz P, Fraser R, Kreiss C, D'Amato M, Crelier G, Boesiger P, Freid M. Role of cholecystokinin in the regulation of liquid gastric emptying and gastric motility in humans: Studies with the CCK antagonist loxiglumide. *Gut* 1997;41:500–504.
111. Mesquita MA, Thompson DG, Troncon LE, D'Amato M, Rovati LC, Barlow J. Effect of cholecystokinin—A receptor blockade on lipid-induced gastric relaxation in humans. *Am J Physiol* 1997;273:G118–123.
112. Trudgill NJ, Hussain FN, Moustafa M, Ajjan R, D'Amato M, Riley SA. The effect of cholecystokinin antagonism on postprandial lower oesophageal sphincter function in asymptomatic volunteers and patients with reflux disease. *Aliment Pharmacol Ther* 2001;15:1357–1364.
113. Persiani S, D'Amato M, Makovec F, Tavares IA, Bishai PM, Rovati LC. Pharmacokinetics of dexloxiglumide after administration of single and repeat oral escalating doses in healthy young males. *Int J Clin Pharmacol Ther* 2002;40:198–206.
114. Varga G. Dexloxiglumide Rotta Research Lab. *Curr Opin Investig Drugs* 2002;3:621–626.
115. Kerwin JF, Nadzan AM, Kopecka H, Lin C W, Miller T, Witte D, Burt S. Hybrid cholecystokinin (CCK) antagonists: New implications in the design and modification of CCK antagonists. *J Med Chem* 1989;32:739–742.
116. Freidinger RM, Whitter WL, Gould NP, Holloway MK, Chang RSL, Lotti VJ. Novel glutamic acid derived cholecystokinin receptor ligands. *J Med Chem* 1990;33:591–595.
117. Kerwin JF, Wagenaar F, Kopecka H, Lin CW, Miller T, Witte D, Stashko M, Nadzan AM. Cholecystokinin antagonists: (R)-tryptophan-based hybrid antagonists of high affinity and selectivity for CCK-A receptors. *J Med Chem* 1991;34:3350–3359.
118. Yamazi Y, Akahane M, Kobayashi M, Kitazawa M, Kurashina Y, Iizuka K. Pharmacological profile of KSG-504, a new cholecystokinin-A-receptor antagonist. *Jpn J Pharmacol* 1993;63:219–225.
119. Hull RA, Shankley NP, Harper EA, Gerkowitch VP, Black JW. 2-Naphthalenesulphonyl-L-aspartyl-(2-phenyl)amide (2-NAP) a selective cholecystokinin CCKA-receptor antagonist. *Br J Pharmacol* 1993;108:734–740.
120. Goursaud AP, Nowak R. 2-NAP, a peripheral CCK-A receptor antagonist, modulates the development of a preference for the mother by the newborn lamb. *Pharmacol Biochem Behav* 2000;67:603–611.
121. Makovec F, Peris W, Revel L, Giovanetti R, Mennuni L, Rovati LC. Structure-antigastrin activity relationships of new (R)-4-benzamido-5-oxopentanoic acid derivatives. *J Med Chem* 1992;35:28–38.
122. Revel L, Ferrari F, Makovec F, Rovati LC, Impicciatore M. Characterization of antagastrin activity *in vivo* of CR 2194, a new R-4-benzamido-5-oxo-pentanoic acid derivative. *Eur J Pharmacol.* 1992;216:217–224.
123. Beltinger J, Hidelbrand P, Howald A, D'Amato M, Beglinger C. Effect of CR 2194, a novel CCK_B/gastrin-receptor antagonist, on gastrin-stimulated acid secretion in man. *Gastroenterology* 1994;106(Suppl): Abstract A50.
124. Beltinger J, Hidelbrand P, Drewe J, Christ A, Hlobil K, Ritz M, D'Amato M, Rovati L, Beglinger C. Effects of spiroglumide, a gastrin receptor antagonist, on acid secretion in humans. *Eur J Clin Invest.* 1999;29:153–159.

125. Makovec F, Peris W, Frigerio S, Giovanetti R, Letari O, Mennuni L, Revel L. Structure-antigastrin activity relationships of new spiroglumide amido acid derivatives. *J Med Chem* 1996;39:135–142.
126. Rovati LC, D'Amato M, Peris W, Revel L, Makovec F. Selectivity and potency of new basic CCK-B antagonists. *Ann NY Acad Sci* 1994;713:395–397.
127. Makovec F, Revel L, Letari O, Mennuni L, Impicciatore M. Characterization of antisecretory and antiulcer activity of CR 2945, a new potent and selective gastrin/CCK_B antagonist. *Eur J Pharmacol* 1999;369:81–90.
128. Revel L, Mennuni L, Garofalo P, Makovec F. CR 2945: A novel CCK_B receptor antagonist with anxiolytic-like activity. *Behav Pharmacol* 1998;9:183–194.
129. Fontana MG, Donato F, Villanacci V, Ghirardi M, Moneghini D, Di Betta E, Salerni B. Inhibitory effect of a gastrin receptor antagonist, CR2945, on 1, 2-dimethylhydrazine-induced colorectal cancer in mice. *Eur Surg Res* 1999;31:406–411.
130. Bock MG, DiPardo RM, Rittle KE, Evans BE, Freidinger RM, Veber DF, Chang RSL, Chen T, Keegan ME, Lotti VJ. Cholecystokinin antagonists. Synthesis of asperlicin analogues with improved potency and water solubility. *J Med Chem* 1986;29:1941–1945.
131. Evans BE, Rittle KE, Bock MG, DiPardo RM, Whitter WL, Gould NP, Lundell GF, Homnick CF, Veber DF, Anderson PS, Chang RSL, Lotti VJ, Cerino DJ, Chen TB, King PJ, Kunkel KA, Springer JP, Hirshfield J. Design of nonpeptidal ligands for peptide receptor: Cholecystokinin antagonists. *J Med Chem* 1987;30:1229–1239.
132. Evans BE, Bock MG, Rittle KE, DiPardo RM, Whitter WL, Veber DF, Anderson PS, Freidinger RM. Design of potent, orally effective, nonpeptidal antagonists of the peptide hormone cholecystokinin. *Proc Natl Acad Sci USA* 1986;83:4918–4922.
133. Evans BE, Rittle KE, Bock MG, DiPardo RM, Freidinger RM, Whitter WL, Lundell GF, Veber DF, Anderson PS, Chang RSL, Lotti VJ, Cerino DJ, Chen TB, Kling PJ, Kunkel KA, Springer JP, Hirshfield J. Methods for drug discovery: Development of potent, selective orally effective cholecystokinin antagonists. *J Med Chem* 1988;31:2235–2246.
134. Pullen RG, Hodgson OJ. Penetration of diazepam and the non-peptide CCK antagonist, L-364,718, into rat brain. *J Pharm Pharmacol* 1987;39:863–864.
135. Woltman TA, Hulce M, Reidelberger RD. Relative blood–brain barrier permeabilities of the cholecystokinin receptor antagonists devazepide and A-65186 in rats. *J Pharm Pharmacol* 1999;51:917–920.
136. Cantor P, Mortensen PE, Myhre J, Gjørup I, Worning H, Stahl E, Survill TT. The effect of the cholecystokinin receptor antagonist MK-329 on meal-stimulated pancreaticobiliary output in humans. *Gastroenterology* 1992;102:1742–1751.
137. Cantor P, Olsen O, Gertz BJ, Gjørup I, Worning H. Inhibition of cholecystokinin-stimulated pancreaticobiliary output in man by the cholecystokinin receptor antagonist MK-329. *Scand J Gastroenterol* 1991;26:627–637.
138. Liddle RA, Gertz BJ, Kanayama S, Beccaria L, Coker LD, Turnbull TA, Morita ET. Effects of a novel cholecystokinin (CCK) receptor antagonist, MK-329, on gall bladder contraction and gastric emptying in humans. Implications for the physiology of CCK. *J Clin Invest* 1989;84:1220–1225.
139. Iversen LL, Dourish CT, Iversen SD. Cholecystokinin receptors: Synthetic antagonists with selectivity for receptor subtypes and possible clinical applications. *Biochem Soc Trans* 1991;19:913–914.
140. Ohlsson B, Axelsson J, Rehfeld JF, Ihse I. Devazepide-induced hyperplasia in the rat liver and bile ducts. *Eur Surg Res* 1996;28:299–305.
141. Satoh Y, Matsuo T, Sogabe H, Itoh H, Tada T, Kinoshita T, Yoshida K, Takaya T. Studies on a novel, potent and orally effective cholecystokinin A antagonist, FK-480. Synthesis and structure-activity relationships of FK-480 and related compounds. *Chem Pharm Bull (Tokyo)* 1994;42:2071–2083.
142. Ito H, Sogabe H, Nakarai T, Sato Y, Tomoi M, Kadowaki M, Matsuo M, Tokoro K, Yoshida K. Pharmacological profile of FK480, a novel cholecystokinin type-A receptor antagonist: Comparison to loxiglumide. *J Pharmacol Exp Ther* 1994;268:571–575.
143. Waldeck H, Benson W, Zeugner H, Wolf KU, Gregory PC, Hamminga D, Van Wijngaarden I. Preparation of 1,7-annelated *N*-(2-oxo-1,4-benzodiazepin-3-yl)-¹H-indole-2-carboxamides as cholecystokinin antagonists. *Ger Offen* 1990; Pat Application: DE 89-3907390.
144. Zabielski R, Lesniewska V, Borlak J, Gregory PC, Kiela P, Pierzynowski SG, Barej W. Effects of intraduodenal administration of tarazepide on pancreatic secretion and duodenal EMG in neonatal calves. *Regul Pept* 1998;78:113–123.
145. Bock MG, DiPardo R, Evans BE, Rittle KE, Whitter WL, Veber DF, Anderson PS, Freidinger RM. Benzodiazepine gastrin and brain cholecystokinin receptor ligands: L-365,260. *J Med Chem* 1989;32:13–16.

146. Lotti VJ, Chang RS. A new potent and selective non-peptide gastrin antagonist and brain cholecystokinin receptor (CCK-B) ligand: L-365,260. *Eur J Pharmacol* 1989;162:273–280.
147. Murphy MG, Sytnik B, Kovacs TO, Mertz H, Ewanik D, Shingo S, Lin JH, Gertz BJ, Walsh JH. The gastrin-receptor antagonist L-365,260 inhibits stimulated acid secretion in humans. *Clin Pharmacol Ther* 1993;54:533–539.
148. Bradwejn J, Koszycki D, Couetoux du Tertre A, van Megen H, den Boer J, Westenberg H. The panicogenic effects of cholecystokinin-tetrapeptide are antagonized by L-365,260, a central cholecystokinin receptor antagonist, in patients with panic disorder. *Arch Gen Psychiatry* 1994;51:486–493.
149. Westenberg HGM, Van Megen HJGM, Den Boer JA. Effects of the cholecystokinin-B antagonist L-365,260 on lactate-induced panic disorder (PD) patients. *Psychopharmacol Bull* 1994;30:644.
150. Lines C, Challenor J, Traub M. Cholecystokinin and anxiety in normal volunteers: An investigation of the anxiogenic properties of pentagastrin and reversal by the cholecystokinin receptor subtype B antagonist L-365,260. *Br J Clin Pharmacol* 1995;39:235–242.
151. Kramer MS, Cutler NR, Ballenger JC, Patterson WM, Mendels J, Chenault A, Shrivastava R, Matzura-Wolfe D, Lines C, Reines S. A placebo-controlled trial of L-365,260, a CCK_B antagonist, in panic disorder. *Biol Psychiatry* 1995;37:462–466.
152. Chen IW, Dorley JM, Ramjit HG, Pitzenberger SM, Lin JH. Physiological disposition and metabolism of L-365,260, a potent antagonist of brain cholecystokinin receptor, in laboratory animals. *Drug Metab Dispos* 1992;20:390–395.
153. Lin JH, Storey DE, Chen IW, Xu X. Improved oral absorption of L-365,260, a poorly soluble drug. *Biopharm Drug Dispos* 1996;17:1–15.
154. Bock MG, DiPardo RM, Evans BE, Rittle KE, Whitter WL, Garsky VM, Gilbert KF, Leighton JL, Carson KL, Mellin EC, Veber DF, Chang RDL, Lotti VJ, Freedman SB, Smith AJ, Patel S, Anderson PS, Freidinger RM. Development of 1,4-benzodiazepine cholecystokinin type B antagonists. *J Med Chem* 1993;36:4276–4292.
155. Chambers MS, Hobbs SC, Fletcher SR, Matass VG, Mitchell PJ, Watt AP, Baker R, Freedman SB, Patel S, Smith AJ. L-708,474: The C5-cyclohexyl analogue of L-365,260, a selective high affinity ligand for the CCK_B/gastrin receptor. *Bioorg Med Chem Lett* 1993;3:1919–1924.
156. Bock MG, DiPardo RM, Mellin EC, Newton RC, Veber DF, Freedman SB, Smith AJ, Patel S, Kemp JA, Marshall GR, Fletcher AE, Chapman KL, Anderson PS, Freidinger RM. Second-generation benzodiazepine CCK-B antagonists. Development of subnanomolar analogs with selectivity and water solubility. *J Med Chem* 1994;37:722–724.
157. Castro JL, Ball RG, Broughton HB, Russell MG, Rathbone D, Watt AP, Baker R, Chapman KL, Fletcher AE, Patel S, Smith AJ, Marshall GR, Ryecroft W, Matassa VG. Controlled modification of acidity in cholecystokinin B receptor antagonists: *N*-(1,4-benzodiazepin-3-yl)-*N'*-[3-(tetrazol-5-ylamino)phenyl]ureas. *J Med Chem* 1996;39:842–849.
158. Showell GA, Bourrain S, Neduvellil JG, Fletcher SR, Baker R, Watt AD, Fletcher AE, Freedman SB, Kemp JA, Marshall GR, Patel S, Smith AJ, Matassa VG. High-affinity and potent, water-soluble 5-amino-1,4-benzodiazepine CCK_B/gastrin receptor antagonists containing a cationic solubilising group. *J Med Chem* 1994;37:719–721.
159. Castro JL, Broughton HB, Russell MG, Rathbone D, Watt AP, Ball RG, Chapman KL, Patel S, Smith AJ, Marshall GR, Matassa VG. 5-(Piperidin-2-yl)- and 5-(homopiperidin-2-yl)-1,4-benzodiazepines: High-affinity, basic ligands for the cholecystokinin-B receptor. *J Med Chem* 1997;40:2491–2501.
160. Showell GA, Bourrain S, Fletcher SR, Neduvellil JG, Fletcher AE, Freedman SB, Patel S, Smith AJ, Marshall GR, Graham MI, Sohal B, Matassa VG. C5-Piperaziny-1,4-benzodiazepines, water-soluble, orally bioavailable CCK_B/gastrin receptor antagonists. *Bioorg Med Chem Lett* 1995;5:3023–3026.
161. Patel S, Smith AJ, Chapman KL, Fletcher AE, Kemp JA, Marshall GR, Hargreaves RJ, Ryecroft W, Iversen LL, Iversen SD, et al. Biological properties of the benzodiazepine amidine derivative L-740,093, a cholecystokinin-B/gastrin receptor antagonist with high affinity *in vitro* and high potency *in vivo*. *Mol Pharmacol* 1994;46:943–948.
162. Selnick HG, Liverton NJ, Baldwin JJ, Butcher JW, Claremon DA, Elliott JM, Freidinger RM, King SA, Libby BE, McIntyre CJ, Pribush DA, Remy DC, Smith GR, Tebben AJ, Jurkiewicz NK, Lynch JJ, Salata JJ, Sanguinetti MC, Siegl PK, Slaughter DE, Vyas K. Class III antiarrhythmic activity *in vivo* by selective blockade of the slowly activating cardiac delayed rectifier potassium current I_{Ks} by (R)-2-(2,4-trifluoromethyl)-*N*-[2-oxo-5-phenyl-1-(2,2,2-trifluoroethyl)-2,3-dihydro-1^H-benzo[e][1,4]diazepin-3-yl]-acetamide. *J Med Chem* 1997;40:3865–3868.
163. Satoh M, Kondoh Y, Okamoto Y, Nishida A, Miyata K, Ohta M, Mase T, Murase K. New 1,4-benzodiazepin-2-one derivatives as gastrin/cholecystokinin-B antagonists. *Chem Pharm Bull* 1995;43:2159–2167.

164. Nishida A, Miyata K, Tsutsumi R, Yuki H, Akuzawa S, Kobayashi A, Kamato T, Ito H, Yamano M, Katuyama Y, et al. Pharmacological profile of (R)-1-[2,3-dihydro-1-(2'-methylphenacyl)-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl]-3-(3-methylphenyl)urea (YM022), a new potent and selective gastrin/cholecystokinin-B receptor antagonist, *in vitro* and *in vivo*. *J Pharmacol Exp Ther* 1994;269:725–731.
165. Yano K, Kajiyama A, Yamazaki S, Matsumura Y, Watanabe K, Yamamoto K. *In vitro* stability and *in vivo* absorption studies of colloidal particles formed from a solid dispersion system. *Chem Pharm Bull (Tokyo)* 1996;44:2309–2313.
166. Nishida A, Takinami Y, Yuki H, Kobayashi A, Akuzawa S, Kamato T, Ito H, Yamano M, Nagakura Y, Miyata K. YM022 [(R)-1-[2,3-dihydro-1-(2'-methylphenacyl)-2-oxo-5-phenyl-1H-1,4-benzodiazepin-3-yl]-3-(3-methylphenyl)urea], a potent and selective gastrin/cholecystokinin-B receptor antagonist, prevents gastric and duodenal lesions in rats. *J Pharmacol Exp Ther* 1994;270:1256–1261.
167. Nishida A, Kobayashi-Uchida A, Akuzawa S, Takinami Y, Shishido T, Kamato T, Ito H, Yamano M, Yuki H, Nagakura Y, et al. Gastrin receptor antagonist YM022 prevents hypersecretion after long-term acid suppression. *Am J Physiol* 1995;269:G699–705.
168. Semple G, Ryder H, Rooker DP, Batt AR, Kendrick DA, Szelke M, Ohta M, Satoh M, Nishida A, Akuzawa S, Miyata K. (3R)-N-(1-(tert-butylcarbonylmethyl)-2,3-dihydro-2-oxo-5-(2-pyridyl)-1H-1,4-benzodiazepin-3-yl)-N'-(3-(methylamino)phenyl)urea (YF476): A potent and orally active gastrin/CCK-B antagonist. *J Med Chem* 1997;40:331–341.
169. Takinami Y, Yuki H, Nishida A, Akuzawa S, Uchida A, Takemoto Y, Ohta M, Satoh M, Semple G, Miyata K. YF476 is a new potent and selective gastrin/cholecystokinin-B receptor antagonist *in vitro* and *in vivo*. *Aliment Pharmacol Ther* 1997;11:113–120.
170. Schmassmann A, Reubi JC. Cholecystokinin-B/gastrin receptors enhance wound healing in the rat gastric mucosa. *J Clin Invest* 2000;106:1021–1029.
171. Kitano M, Norlen P, Ding XQ, Nakamura S, Hakanson R. Long-lasting cholecystokinin(2) receptor blockade after a single subcutaneous injection of YF476 or YM022. *Br J Pharmacol* 2000;130:699–705.
172. Kawanishi Y, Ishihara S, Tsushima T, Seno K, Miyagoshi M, Hagishita S, Ishikawa M, Shima N, Shimamura M, Ishihara Y. Design, synthesis and pharmacological evaluation of dual histamine H₂ and gastrin receptor antagonists. *Bioorg Med Chem Lett* 1996;6:1421–1426.
173. Kawanishi Y, Ishihara S, Tsushima T, Seno K, Miyagoshi M, Hagishita S, Ishikawa M, Shima N, Shimamura M, Ishihara Y. Synthesis and pharmacological evaluation of highly potent dual histamine H₂ and gastrin antagonists. *Bioorg Med Chem Lett* 1996;6:1427–1430.
174. Kawanishi Y, Ishihara S, Takahashi K, Tsushima T, Hagishita S, Ishikawa M, Ishihara Y. Synthesis and biological evaluation of a new reversely linked type of dual histamine H₂ and gastrin receptor antagonist. *Chem Pharm Bull* 1997;45:116–124.
175. Satoh Y, Tabuchi S, Mitsui H. Design of dual CCK-A and CCK-B receptor antagonists. *Drugs of the Future* 1997;22:117–1131.
176. Tabuchi S, Nakanishi I, Satoh Y. Relationship between dihedral angles of N1 and C9 substituents in 1,4-benzodiazepines and dual cholecystokinin-A and -B antagonistic activities. *Bioorg Med Chem Lett* 1998;8:1449–1454.
177. Tabuchi S, Ito H, Sogabe H, Kuno M, Kinoshita T, Katumi I, Yamamoto N, Mitsui H, Satoh Y. Dual CCK-A and CCK-B receptor antagonists (II). Preparation and structure activity relationships of 5-alkyl-9-methyl-1,4-benzodiazepines and discovery of FR208419. *Chem Pharm Bull* 2000;48:1–15.
178. Bock MG, DiPardo RM, Veber DF, Chang RSL, Lotti VJ, Freedman SB, Freidinger RM. Benzolactams as non-peptide cholecystokinin receptor ligands. *Bioorg Med Chem Lett* 1993;3:871–874.
179. Lowe JA 3rd, Hageman DL, Drozda SE, McLean S, Bryce DK, Crawford RT, Zorn S, Morrone J, Bordner J. 5-Phenyl-3-ureidobenzazepin-2-ones as cholecystokinin-B receptor antagonists. *J Med Chem* 1994;37:3789–3811.
180. Lowe JA, Drozda AE, McLean S, Bryce DK, Crawford RT. A water soluble benzazepine cholecystokinin-B receptor antagonist. *Bioorg Med Chem Lett* 1995;5:1933–1936.
181. Aquino CJ, Armour DR, Berman JM, Birkemo LS, Carr RA, Croom DK, Dezube M, Dougherty RW Jr, Ervin GN, Grizzle MK, Head JE, Hirst GC, James MK, Johnson MF, Miller LJ, Queen KL, Rimele TJ, Smith DN, Sugg EE. Discovery of 1,5-benzodiazepines with peripheral cholecystokinin (CCK-A) receptor agonist activity. 1. Optimization of the agonist "trigger". *J Med Chem* 1996;39:562–569.
182. Ursini A, Capelli AM, Carr RA, Cassara P, Corsi M, Curcuruto O, Curotto G, Dal Cin M, Davalli S, Donati D, Feriani A, Finch H, Finizia G, Gaviraghi G, Marien M, Pentassuglia G, Polinelli S, Ratti E, Reggiani AM, Tarzia G, Tedesco G, Tranquillini ME, Trist DG, Van Amsterdam FT, Reggiani A. Synthesis and SAR of new 5-phenyl-3-ureido-1,5-benzodiazepines as cholecystokinin-B receptor antagonists. *J Med Chem* 2000;43:3596–3613.

183. Reggiani A, Gerrard PA, Maraia G, Melotto S, Ratti E, Gaviraghi G, Trist DG. Anxiolytic activity of GV150013, a new potent and selective cholecystokinin-B (CCK-B) antagonist. *Br J Pharmacol* 1995; 114(Proceedings Suppl):92P.
184. Crespi F. Cholecystokinin-B (CCK-B) receptor antagonists improve "aged" sleep: A new class of sleep modulators? *Methods Find Exp Clin Pharmacol* 1999;21:31-38.
185. Bailey N, Box PC, Carr RAE, Cooke JWB, Evans B, Finch H, Head JE, Pass M, Shah P, Wheatcroft JR. Novel 1,5-benzodiazepindione gastrin/CCK_B antagonists. *Bioorg Med Chem Lett* 1997;7:281-286.
186. Hagishita A, Seno K, Kamata S, Haga N, Ishihara Y, Ishikawa M, Shimamura M. Potent and subtype-selective CCK-B/gastrin receptor antagonists: 2,4-Dioxo-1,5-benzodiazepines with a plane of symmetry. *Bioorg Med Chem* 1997;5:1433-1446.
187. Bohme GA, Bertrand P, Guyon C, Capet M, Pendley C, Stutzmann JM, Doble A, Dubroeuq MC, Martin G, Blanchard JC. The ureidoacetamides, a novel family of non-peptide CCK-B/gastrin antagonists. *Ann N Y Acad Sci* 1994;713:118-120.
188. Bertrand P, Bohme GA, Durieux C, Guyon C, Capet M, Jeantaud B, Boudeau P, Ducos B, Pendley CE, Martin GE, et al. Pharmacological properties of ureido-acetamides, new potent and selective non-peptide CCKB/gastrin receptor antagonists. *Eur J Pharmacol* 1994;262:233-245.
189. Pendley CE, Fitzpatrick LR, Capolino AJ, Davis MA, Esterline NJ, Jakubowska A, Bertrand P, Guyon C, Dubroeuq MC, Martin GE. RP 73870, a gastrin/cholecystokinin-B receptor antagonist with potent anti-ulcer activity in the rat. *J Pharmacol Exp Ther* 1995;273:1015-1022.
190. Hagishita S, Murakami Y, Seno K, Kamata S, Haga N, Konoike T, Kanda Y, Kiyama R, Shiota T, Ishihara Y, Ishikawa M, Shimamura M, Abe K, Yoshimura K. Synthesis and pharmacological properties of ureidomethylcarbonylphenylketone derivatives. A new potent and subtype-selective nonpeptide CCK-B/gastrin receptor antagonist, S-0509. *Bioorg Med Chem* 1997;5:1695-1714.
191. Takeda Y, Kawagoe K, Yokomizo A, Yokomizo Y, Hosokami T, Ogihara Y, Honda Y, Yokohama S. Synthesis of phenoxyacetic acid derivatives as highly potent antagonists of gastrin/cholecystokinin-B receptors. *Chem Pharm Bull (Tokyo)* 1998;46:434-444.
192. Takeda Y, Kawagoe K, Yokomizo A, Yokomizo Y, Hosokami T, Shimoto Y, Tabuchi Y, Ogihara Y, Otsubo R, Honda Y, Yokohama S. Synthesis of phenoxyacetic acid derivatives as highly potent antagonists of gastrin/cholecystokinin-B receptors. II. *Chem Pharm Bull (Tokyo)* 1998;46:951-961.
193. Takeda Y, Kawagoe K, Yokomizo A, Yokomizo Y, Hosokami T, Shimoto Y, Tabuchi Y, Ogihara Y, Honda Y, Kawarabayashi K, Iseri M, Yokohama S. Synthesis of phenoxyacetic acid derivatives as highly potent antagonists of gastrin/cholecystokinin-B receptors. III. *Chem Pharm Bull (Tokyo)* 1999;47:755-771.
194. Yu MJ, Thrasher KJ, McCowan JR, Mason NR, Mendelsohn LG. Quinazolinone cholecystokinin-B receptor ligands. *J Med Chem* 1991;34:1505-1508.
195. Rasmussen K, Yu MJ, Czachura JF. Quinazolinone cholecystokinin (CCK)-B antagonists decrease midbrain dopamine unit activity. *Synapse* 1994;17:278-282.
196. Yu MJ, McCowan JR, Mason NR, Deeter JB, Mendelsohn LG. Synthesis and X-ray crystallographic analysis of quinazolinone cholecystokinin/gastrin receptor ligands. *J Med Chem* 1992;35:2534-2542.
197. Padia JK, Chilvers H, Daum P, Pinnock R, Suman-Chauhan N, Webdale L, Trivedi BK. Design and synthesis of novel nonpeptide CCK-B receptor antagonists. *Bioorg Med Chem Lett* 1997;7:805-810.
198. Padia JK, Field M, Hinton J, Meecham K, Pablo J, Pinnock R, Roth BD, Singh L, Suman-Chauhan N, Trivedi BK, Webdale L. Novel nonpeptide CCK-B antagonists: Design and development of quinazolinone derivatives as potent, selective, and orally active CCK-B antagonists. *J Med Chem* 1998;41:1042-1049.
199. Horwell DC, Beeby A, Clark CR, Hughes J. Synthesis and binding affinities of analogues of cholecystokinin-(30-33) as probes for central nervous system cholecystokinin receptors. *J Med Chem* 1987;30:729-732.
200. Horwell DC, Birchmore B, Boden PR, Higginbottom M, Ping Ho Y, Hughes J, Hunter JC, Richardson RS. α -Methyl tryptophanylphenylalanines and their aryethylamine "dipeptoid" analogues of the tetrapeptide cholecystokinin (30-33). *Eur J Med Chem* 1990;25:53-60.
201. Augelli-Szafran CE, Purchase TS, Roth BD, Tait B, Trivedi BK, Wilson M, Suman-Chauhan N, Webdale L. SAR study of the indole moiety of CI-988, a potent and selective CCK-B antagonist. *Bioorg Med Chem Lett* 1997;7:2009-2014.
202. Horwell DC, Hughes J, Hunter JC, Pritchard MC, Richardson RS, Roberts E, Woodruff GN. Rationally designed "dipeptoid" analogues of CCK. α -Methyltryptophan derivatives as highly selective and orally active gastrin and CCK-B antagonists with potent anxiolytic properties. *J Med Chem* 1991;34:404-414.
203. Drysdale MJ, Pritchard MC, Horwell DC. Rationally designed "dipeptoid" analogues of CCK. Acid mimics of the potent and selective non-peptide CCK-B receptor antagonist CI-988. *J Med Chem* 1992;35:2573-2581.

204. Hughes J, Boden P, Costall B, Domeney A, Kelly E, Horwell DC, Hunter JC, Pinnock RD, Woodruff GN. Development of a class of selective cholecystokinin type B receptor antagonists having potent anxiolytic activity. *Proc Natl Acad Sci USA* 1990;87:6728–6732.
205. Adams JB, Pyke RE, Costa J, Cutler NR, Schweizer E, Wilcox CS, Wisselink PG, Greiner M, Pierce MW, Pande AC. A double-blind, placebo-controlled study of a CCK-B receptor antagonist, CI-988, in patients with generalized anxiety disorder. *J Clin Psychopharmacol* 1995;15:428–434.
206. van Megen HJ, Westenberg HG, den Boer JA, Slaap B, van Es-Radhakishun F, Pande AC. The cholecystokinin-B receptor antagonist CI-988 failed to affect CCK-4 induced symptoms in panic disorder patients. *Psychopharmacology* 1997;129:243–248.
207. Goddard AW, Woods SW, Money R, Pande AC, Charney DS, Goodman WK, Heninger GR, Price LH. Effects of the CCK(B) antagonist CI-988 on responses to mCPP in generalized anxiety disorder. *Psychiatry Res* 1999;85:225–240.
208. Pande AC, Greiner M, Adams JB, Lydiard RB, Pierce MW. Placebo-controlled trial of the CCK-B antagonist, CI-988, in panic disorder. *Biol Psychiatry* 1999;46:860–862.
209. Cowley DS, Adams JB, Pyke RE, Cook J, Zaccharias P, Wingerson D, Roy-Byrne PP. Effect of CI-988, a cholecystokinin-B receptor antagonist, on lactate-induced panic. *Biol Psychiatry* 1996;40:550–552.
210. Bradwejn J, Koszycki D, Paradis M, Reece P, Hinton J, Sedman A. Effect of CI-988 on cholecystokinin tetrapeptide-induced panic symptoms in healthy volunteers. *Biol Psychiatry* 1995;38:742–746.
211. Ding X-Q, Chen D, Håkanson R. Cholecystokinin-B/gastrin receptor ligands of the dipeptoid series act as agonists on histidine decarboxylase in rat stomach enterochromaffinlike cells. *Pharmacol Toxicol* 1995;76(Suppl IV):81.
212. Marvik R, Sandvik AK, Waldum HL. Evaluation of biologic gastrin activity of compound CI-988 in the isolated, vascularly perfused rat stomach. *Scand J Gastroenterol* 1997;32:519–522.
213. Patel S, Chapman KL, Smith AJ, Bailey I, Freedman SB. Are radioligand antagonist/agonist binding ratios in rat pancreas predictive of functional efficacy of cholecystokinin receptor agonists and antagonists? *Regul Pept* 1996;65:29–35.
214. Coudore-Civiale MA, Courteix C, Boucher M, Meen M, Fialip J, Eschalier A, Ardid D. Potentiation of morphine and clomipramine analgesia by cholecystokinin-B antagonist CI-988 in diabetic rats. *Neurosci Lett* 2000;286:37–40.
215. Coudore-Civiale MA, Courteix C, Fialip J, Boucher M, Eschalier A. Spinal effect of the cholecystokinin-B receptor antagonist CI-988 on hyperalgesia, allodynia and morphine-induced analgesia in diabetic and mononeuropathic rats. *Pain* 2000;88:15–22.
216. Trivedi BK, Padia JK, Holmes A, Rose S, Wright DS, Hinton JP, Pritchard MC, Eden JM, Kneen C, Webdale L, Suman-Chauhan N, Boden P, Singh L, Field MJ, Hill D. Second generation "peptoid" CCK-B receptor antagonists: Identification and development of *N*-(adamantylloxycarbonyl)- α -methyl-(*R*)-tryptophan derivative (CI-1015) with an improved pharmacokinetic profile. *J Med Chem* 1998;41:38–45.
217. Trivedi BK, Hinton JP. CI-1015. An orally active CCK-B receptor antagonist with an improved pharmacokinetic profile. *Pharm Biotechnol* 1998;11:481–505.
218. Fincham CI, Higginbottom M, Hill DR, Horwell DC, O'Toole JC, Ratcliffe GS, Rees DC, Roberts E. Amide bond replacements incorporated into CCK-B selective "dipeptoids". *J Med Chem* 1992;35:1472–1484.
219. Boden PR, Higginbottom M, Hill DR, Horwell DC, Hughes J, Rees DC, Roberts E, Singh L, Suman-Chauhan N, Woodruff GN. Cholecystokinin dipeptoid antagonists: Design, synthesis, and anxiolytic profile of some novel CCK-A and CCK-B selective and "mixed" CCK-A/CCK-B antagonists. *J Med Chem* 1993;36:552–565.
220. Augelli-Szafran CE, Horwell DC, Kneen C, Ortwin DF, Pritchard MC, Purchase TS, Roth BD, Trivedi BK, Hill D, Suman-Chauhan N, Webdale L. Cholecystokinin B antagonists. Synthesis and quantitative structure-activity relationships of a series of C-terminal analogues of CI-988. *Bioorg Med Chem* 1996;4:1733–1745.
221. Hughes J, Dockray GJ, Hill D, Garcia L, Pritchard MC, Forster E, Toescu E, Woodruff G, Horwell DC. Characterization of novel peptoid agonists for the CCK-A receptor. *Regul Pept* 1996;65:15–21.
222. Fincham CI, Horwell DC, Ratcliffe GS, Rees DC. The use of a proline ring as a conformational restraint in CCK-B receptor "dipeptoids". *Bioorg Med Chem Lett* 1992;2:403–406.
223. Higginbottom M, Hill DR, Horwell DC, Mostafai E, Suman-Chauhan N, Roberts E. Conformationally restricted analogues of the potent CCK-B antagonist CI-988. *Bioorg Med Chem* 1993;1:209–217.
224. Bellier B, McCort-Tranchepain I, Ducos B, Danascimento S, Meudal H, Noble F, Garbay C, Roques BP. Synthesis and biological properties of new constrained CCK-B antagonists: Discrimination of two affinity states of the CCK-B receptor on transfected CHO cells. *J Med Chem* 1997;40:3947–3956.

225. Eden JM, Horwell DC, Pritchard MC. α , β -didehydrotryptophan as a surrogate for α -methyltryptophan in CCK 'peptoids' related to CI-988. *Bioorg Med Chem Lett* 1993;3:989–992.
226. De la Figuera N, Martín-Martínez M, Herranz R, García-López MT, Latorre M, Cenarruzabeitia E, Del Río J, González-Muñiz R. Highly constrained dipeptoid analogues containing a type II' β -type mimic as novel and selective CCK-A receptor ligands. *Bioorg Med Chem Lett* 1999;9:43–48.
227. Martín-Martínez M, De La Figuera N, Latorre M, Herranz R, García-López MT, Cenarruzabeitia E, Del Río J, González-Muñiz R. β -Turned dipeptoids as potent and selective CCK₁ receptor antagonists. *J Med Chem* 2000;43:3770–3777.
228. Kalindjian SB, Bodkin MJ, Buck IM, Dunstone DJ, Low CM, McDonald IM, Pether MJ, Steel KI. A new class of non-peptidic cholecystokinin-B/gastrin receptor antagonists based on dibenzobicyclo[2.2.2]octane. *J Med Chem* 1994;37:3671–3673.
229. Kalindjian SB, Buck IM, Cushnir JR, Dunstone DJ, Hudson ML, Low CM, McDonald IM, Pether MJ, Steel KI, Tozer MJ. Improving the affinity and selectivity of a nonpeptide series of cholecystokinin-B/gastrin receptor antagonists based on the dibenzobicyclo[2.2.2]octane skeleton. *J Med Chem* 1995;38:4294–4302.
230. Kalindjian SB, Buck IM, Davies JM, Dunstone DJ, Hudson ML, Low CM, McDonald IM, Pether MJ, Steel KI, Tozer MJ, Vinter JG. Non-peptide cholecystokinin-B/gastrin receptor antagonists based on bicyclic, heteroaromatic skeletons. *J Med Chem* 1996;39:1806–1815.
231. Ding XQ, Lindstrom E, Hakanson R. Evaluation of three novel cholecystokinin-B/gastrin receptor antagonists: A study of their effects on rat stomach enterochromaffin-like cell activity. *Pharmacol Toxicol* 1997;81:232–237.
232. Lindstrom E, Björkqvist M, Hakanson R. Pharmacological analysis of CCK₂ receptor antagonists using isolated rat stomach ECL cells. *Br J Pharmacol* 1999;127:530–536.
233. Hakanson R, Ding XQ, Norlen P, Lindstrom E. CCK₂ receptor antagonists: Pharmacological tools to study the gastrin-ECL cell-parietal cell axis. *Regul Pept* 1999;80:1–12.
234. Gibson (nee Thomas) SE, Guillo N, Jones JO, Buck IM, Kalindjian SB, Roberts S, Tozer MJ. CCK₂ receptor antagonists containing the conformationally constrained phenylalanine derivatives, including the new amino acid Xic. *Eur J Med Chem* 2002;37:379–389.
235. McDonald IM, Dunstone DJ, Kalindjian SB, Linney ID, Low CM, Pether MJ, Steel KI, Tozer MJ, Vinter JG. 2,7-Dioxo-2,3,4,5,6,7-hexahydro-1*H*-benzo[*h*][1,4]diazonine as a new template for the design of CCK₂ receptor antagonists. *J Med Chem* 2000;43:3518–3529.
236. Kalindjian SB, Dunstone DJ, Low CM, Pether MJ, Roberts SP, Tozer MJ, Watt GF, Shankley NP. Nonpeptide cholecystokinin-2 receptor agonists. *J Med Chem* 2001;44:1125–1133.
237. González-Muñiz R, Domínguez MJ, Martín-Martínez M, Herranz R, García-López MT, Barber A, Ballaz S, Del Río J. CCK-4 restricted analogues containing a 3-oxoindolizidine skeleton. *Bioorg Med Chem Lett* 1996;6:967–972.
238. Martín-Martínez M, Bartolomé-Nebreda JM, Gómez-Monterrey I, González-Muñiz R, García-López MT, Ballaz S, Barber A, Fortuño A, Del Río J, Herranz R. Synthesis and stereochemical structure-activity relationships of 1,3-dioxoperhydropyrido[1,2-*c*]-pyrimidine derivatives: Potent and selective cholecystokinin-A receptor antagonists. *J Med Chem* 1997;40:3402–3407.
239. Ballaz S, Barber A, Fortuño A, Del Río J, Martín-Martínez M, Gómez-Monterrey I, Herranz R, González-Muñiz R, García-López MT. Pharmacological evaluation of IQM-95,333, a highly selective CCK_A receptor antagonist with anxiolytic-like activity in animal models. *Br J Pharmacol* 1997;121:759–767.
240. Woodruff GN, Hill DR, Boden P, Pinnock R, Singh L, Hughes J. Functional role of brain CCK receptors. *Neuropeptides* 1991;19(Suppl):45–56.
241. Hendrie CA, Dourish CT. Anxiolytic profile of the cholecystokinin antagonist devazepide in mice. *Br J Pharmacol* 1990;99:138P.
242. Hendrie CA, Neill JC, Shepherd JK, Dourish CT. The effects of CCKA and CCKB antagonists on activity in the black/white exploration model of anxiety in mice. *Physiol Behav* 1993;54:689–693.
243. Ravard S, Dourish CT. Cholecystokinin and anxiety. *Trends Pharmacol Sci* 1990;11:271–273.
244. Bartolomé-Nebreda JM, Gómez-Monterrey I, García-López MT, González-Muñiz R, Martín-Martínez M, Ballaz S, Cenarruzabeitia E, Latorre M, Del Río J, Herranz R. 5-(Tryptophyl)amino-1,3-dioxoperhydropyrido[1,2-*c*]pyrimidine-based potent and selective CCK₁ receptor antagonists: Structural modifications at the tryptophan domain. *J Med Chem* 1999;42:4659–4668.
245. Bartolomé-Nebreda JM, Patiño-Molina R, Martín-Martínez M, Gómez-Monterrey I, García-López MT, González-Muñiz R, Cenarruzabeitia E, Latorre M, Del Río J, Herranz R. 5-(Tryptophyl)amino-1,3-dioxoperhydropyrido[1,2-*c*]pyrimidine-based potent and selective CCK₁ receptor antagonists: Structure-activity relationship studies on the substituent at N2-position. *J Med Chem* 2001;44:2219–2228.

246. Bartolomé-Nebreda JM, García-López MT, González-Muñiz R, Cenarruzabeitia E, Latorre M, Del Río J, Herranz R. 5-(Tryptophyl)amino-1,3-dioxoperhydropyrido[1,2-c]pyrimidine-based potent and selective CCK₁ receptor antagonists: Structure-activity relationship studies on the central 1,3-dioxoperhydropyrido[1,2-c]pyrimidine scaffold. *J Med Chem* 2001;44:4196–4206.
247. Muñoz-Ruiz P, García-López MT, Cenarruzabeitia E, Del Río J, Herranz R. 1,3-Dioxoperhydropyrido[1,2-c]pyrimidine-based potent and selective CCK receptor antagonists: Reversal of the CCK₁ subtype receptor selectivity towards CCK₂ receptors. Personal communication to the XVIIth International Symposium on Medicinal Chemistry, Barcelona, September 2002.
248. Howbert JJ, Lobb KL, Britton TC, Mason NR, Bruns RF. Diphenylpyrazolidinone and benzodiazepine cholecystokinin antagonists: A case of convergent evolution in medicinal chemistry. *Bioorg Med Chem Lett* 1993;5:875–880.
249. Ding XQ, Hakanson R. Evaluation of the specificity and potency of a series of cholecystokinin-B/gastrin receptor antagonists *in vivo*. *Pharmacol Toxicol* 1996;79:124–130.
250. Helton DR, Berger JE, Czachura JF, Rasmussen K, Kallman MJ. Central nervous system characterization of the new cholecystokininB antagonist LY288513. *Pharmacol Biochem Behav* 1996;53:493–502.
251. Izumi T, Inoue T, Tsuchiya K, Hashimoto S, Ohmori T, Koyama T. Effect of the selective CCKB receptor antagonist LY288513 on conditioned fear stress in rats. *Eur J Pharmacol* 1996;300:25–31.
252. Adamec RE, Shallow T, Budgell J. Blockade of CCK(B) but not CCK(A) receptors before and after the stress of predator exposure prevents lasting increases in anxiety-like behavior: Implications for anxiety associated with posttraumatic stress disorder. *Behav Neurosci* 1997;111:435–449.
253. Rasmussen K, Czachura JF, Kallman MJ, Helton DR. The CCK-B antagonist LY288513 blocks the effects of nicotine withdrawal on auditory startle. *Neuroreport* 1996;7:1050–1052.
254. Rasmussen K, Helton DR, Berger JE, Searce E. The CCK-B antagonist LY288513 blocks diazepam-withdrawal-induced increases in auditory startle response. *Ann N Y Acad Sci* 1994;713:374–376.
255. Flynn DL, Villamil CI, Becker DP, Gullikson GW, Moumni C, Yang D-C. 1,3,4-trisubstituted pyrrolidinones as scaffolds for construction of peptidomimetic cholecystokinin antagonists. *Bioorg Med Chem Lett* 1992;2:1251–1256.
256. Lowe JA III, Qian W, Scott PJ, McLean S, Bryce DK, Crawford RT, Bordner J. 5,7-Diphenyl-3-ureidohexahydroazepin-2-ones as Cholecystokinin-B receptor ligands. *Bioorg Med Chem Lett* 1994;4:2877–2882.
257. Baba A, Hosshino E, Onoma M, Esaki T, Takeda K. Pharmacological profiles of AG-041R, a novel selective gastrin/CCK_B antagonist. *Jpn J Pharmacol* 1995;67(Suppl 1): Abstract P2–119.
258. Baba A, Hosshino E, Onoma M, Esaki T, Takeda K. Pharmacological profiles of AG-041R, a novel selective gastrin/CCK_B antagonist. *Gastroenterology* 1995;108(Suppl):A950.
259. Chibā T, Kinoshita Y, Sawada M, Kishi K, Baba A, Hoshino E. The role of endogenous gastrin in the development of enterochromaffin-like cell carcinoid tumors in *Mastomys natalensis*: A study with the specific gastrin receptor antagonist AG-041R. *Yale J Biol Med* 1998;71:247–255.
260. Kitamura H, Kato A, Esaki T. AG-041R, a novel indoline-2-one derivative, induces systemic cartilage hyperplasia in rats. *Eur J Pharmacol* 2001;418:225–230.
261. Ochi M, Kawasaki K, Kataoka H, Uchio Y, Nishi H. AG-041R, a gastrin/CCK-B antagonist, stimulates chondrocyte proliferation and metabolism *in vitro*. *Biochem Biophys Res Commun* 2001;283:1118–1123.
262. Nakanishi T, Kawasaki K, Uchio Y, Kataoka H, Terashima M, Ochi M. AG-041R, a cholecystokinin-B/gastrin receptor antagonist, stimulates the repair of osteochondral defect in rabbit model. *Eur J Pharmacol* 2002;439:135–140.
263. Taniguchi H, Yazaki N, Endo T, Nagasaki M. Pharmacological profile of T-0632, a novel potent and selective CCKA receptor antagonist, *in vitro*. *Eur J Pharmacol* 1996;304:147–154.
264. Taniguchi H, Yazaki N, Yomota E, Shikano T, Endo T, Nagasaki M. Pharmacological profile of T-0632, a novel potent and selective CCKA receptor antagonist, *in vivo*. *Eur J Pharmacol* 1996;312:227–233.
265. Taniguchi H, Yomota E, Kume E, Shikano T, Endo T, Nagasaki M. Effect of T-0632, a cholecystokininA receptor antagonist, on experimental acute pancreatitis. *Jpn J Pharmacol* 1997;73:105–112.
266. Niederau C, Grendell JH. Role of cholecystokinin in the development and progression of acute pancreatitis and the potential of therapeutic application of cholecystokinin receptor antagonists. *Digestion* 1999; 60Suppl 1:69–74.
267. Gully D, Frehel D, Marcy C, Spinazze A, Lespy L, Neliat G, Maffrand JP, Le Fur G. Peripheral biological activity of SR 27897: A new potent non-peptide antagonist of CCKA receptors. *Eur J Pharmacol* 1993; 232:13–19.
268. Poncelet M, Arnone M, Heaulme M, Gonalons N, Gueudet C, Santucci V, Thurneyssen O, Keane P, Gully D, Le Fur G, et al. Neurobehavioural effects of SR 27897, a selective cholecystokinin type A (CCK-A) receptor antagonist. *Naunyn Schmiedeberg Arch Pharmacol* 1993;348:102–107.

269. Le Drean G, Le Huerou-Luron I, Gestin M, Rome V, Bernard C, Chayvialle JA, Fourmy D, Guilloteau P. Pancreatic secretory response to feeding in the calf: CCK-A receptors, but not CCK-B/gastrin receptors are involved. *Can J Physiol Pharmacol* 2000;78:813–819.
270. González AA, Farre R, Mones J, Capella G, Clave P. Pharmacological and molecular characterization of muscular cholecystokinin receptors in the human lower oesophageal sphincter. *Neurogastroenterol Motil* 2000;12:539–546.
271. Bunting MW, Beart PM, Widdop RE. Effects of the CCK(A) receptor antagonists SR 27897B and PD140548 on baroreflex function in conscious rats. *Eur J Pharmacol* 1997;337:35–39.
272. Matto V, Harro J, Allikmets L. The effects of cholecystokinin A and B receptor antagonists on exploratory behaviour in the elevated zero-maze in rat. *Neuropharmacology* 1997;36:389–396.
273. Kreiss C, Schwizer W, Borovicka J, Jansen JB, Bouloux C, Pignol R, Bischof Delaloye A, Fried M. Effect of linitript, a new CCK-A receptor antagonist, on gastric emptying of a solid-liquid meal in humans. *Regul Pept* 1998;74:143–149.
274. Gouldson P, Legoux P, Carillon C, Delpech B, Le Fur G, Ferrara P, Shire D. Contrasting roles of leu³⁵⁶ in the human CCK₁ receptor for antagonist SR 27897 and agonist SR 146131 binding. *Eur J Pharmacol* 1999;383:339–346.
275. Gigoux V, Maigret B, Escrieut C, Silvente-Poirot S, Bouisson M, Fehrentz JA, Moroder L, Gully D, Martinez J, Vaysse N, Fourmy AD. Arginine 197 of the cholecystokinin-A receptor binding site interacts with the sulfate of the peptide agonist cholecystokinin. *Protein Sci* 1999;8:2347–2354.
276. Gouldson P, Legoux P, Carillon C, Dumont X, Le Fur G, Ferrara P, Shire D. Essential role of extracellular charged residues of the human CCK₁ receptor for interactions with SR 146131, SR 27897 and CCK-8S. *Eur J Pharmacol* 2000;389:115–124.
277. Gouldson P, Legoux P, Carillon C, Delpech B, Le Fur G, Ferrara P, Shire D. The agonist SR 146131 and the antagonist SR 27897 occupy different sites on the human CCK₁ receptor. *Eur J Pharmacol* 2000;400:185–194.
278. Akiyama T, Tachibana I, Hirohata Y, Shirohara H, Yamamoto M, Otsuki M. Pharmacological profile of TP-680, a new cholecystokininA receptor antagonist. *Br J Pharmacol* 1996;117:1558–1564.
279. Tachibana I, Kanagawa K, Yamamoto Y, Otsuki M. Pharmacological profile of a new serine derivative cholecystokinin receptor antagonist TP-680 on pancreatic, biliary and gastric function. *J Pharmacol Exp Ther* 1996;279:1404–1412.
280. Ohtsuka T, Kotaki H, Nakayama N, Itezono Y, Shimma N, Kudoh T, Kuwahara T, Arisawa M, Yokose K, Seto H. Tetronothiodin, a novel cholecystokinin type-B receptor antagonist produced by *Streptomyces* sp. NR0489. II. Isolation, characterization and biological activities. *J Antibiot (Tokyo)* 1993;46:11–17.
281. Ohtsuka T, Nakayama N, Itezono Y, Shimma N, Kuwahara T, Yokose K, Seto H. Tetronothiodin, a novel cholecystokinin type-B receptor antagonist produced by *Streptomyces* sp. NR0489. III. Structural elucidation. *J Antibiot (Tokyo)* 1993;46:18–24.
282. Dugé V, Sebret A, Beslot F, Matsui T, Roques BP. Behavioral profile of CCK2 receptor-deficient mice. *Neuropsychopharmacology* 2001;25:690–698.

Rosario Herranz received her PhD degree in chemistry from the Autonomia University of Madrid, Spain, in 1980. After working as a postdoctoral student in the Medicinal Chemistry Institute of Madrid under the guidance of Professor M. Teresa García-López for 3 years, she joined the Research Department of the pharmaceutical company Antibióticos S.A., as staff researcher until 1986. At this year, she returned to the Medicinal Chemistry Institute, where she is currently Scientific Researcher. Her research interests have evolved from the area of nucleoside analogues, with antitumoral and antiviral activity towards peptide analogues and peptidomimetics with action upon the CNS. In this field, her work has firstly been focused on analgesia, and more recently on the search of non-peptide ligands for cholecystokinin receptors and heterocyclic-derived ligands for glutamate receptors, with a potential application to neuroprotection.